

10-16-00

A

10/13/00

10/13/00

10/13/00

**DIVISION - CONTINUATION - CONTINUATION-IN-PART
APPLICATION TRANSMITTAL FORM**

Attorney Docket No.:

A-357B

Anticipated Classification Of This Application:
Class 530 Subclass

Prior Application:
Examiner S. Gucker

Art Unit 1647

To the Assistant Commissioner for Patents:

This is a request for filing a ☐ continuation ☒ divisional ☐ continuation-in-part application, under 37 CFR 1.53(b),
of pending prior application Serial No. 08/535,681 filed on September 28, 1995,
of Shaw-Fen Sylvia Hu

for **TRUNCATED GLIAL CELL LINE-DERIVED NEUROTROPHIC FACTOR**

For CONTINUATION or DIVISIONAL APPLNS only: The entire disclosure of the prior application, from which an oath or declaration is supplied under Box 1b, below, is considered a part of the disclosure of the accompanying continuation or divisional application and is hereby incorporated by reference. The incorporation can only be relied upon when a portion has been inadvertently omitted from the submitted application parts.

1. ☒ Transmitted herewith are:
☒ 82 pages of specification, 8 pages of claim(s) and 1 page of abstract, totaling 91 pages.
☒ 9 sheet(s) of drawings.
☒ 2 pages of Oath or Declaration by the applicant(s):
☐ a. Newly executed (original or copy)
☒ b. Copy from a prior application (37 CFR 1.63(d)) (for continuation/divisional applns. only)
☒ 22 pages of Sequence Listing

2. ☒ The filing fee is calculated below:

For	Number Filed		Number Extra		Rate		Fee
Total Claims	12	- 20 =	0	x	\$18.00	=	\$ 0.00
Independent Claims	2	- 3 =	0	x	\$80.00	=	0.00
Multiple Dependent Claims	1			+	\$270.00	=	270.00
Basic Fee					\$710.00	=	710.00
Total Filing Fee							\$ 980.00

3. ☒ Please charge Deposit Account No. 01-0519, in the name of Amgen Inc., in the amount of \$ 980.00. An original and one copy are enclosed.
4. ☒ Throughout the prosecution of this application, if any extension of time is necessary, please consider this a request therefor.
5. ☒ The Commissioner is hereby authorized to charge any additional filing fees which may be required by the accompanying application, any additional fees which may be required during pendency of this application as required by 37 CFR 1.16 or 1.17, or credit any overpayment to Deposit Account No. 01-0519 throughout the prosecution of this application.
6. ☒ Cancel in this application original claims 1-29 AND 31-44 of the prior application before calculating the filing fee. (At least one original independent claim must be retained for filing purposes.)

EXPRESS MAIL CERTIFICATE

"Express Mail" mail labeling number: EL90069268US

Date of Deposit: October 13, 2000

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to Box Patent Application, Assistant Commissioner for Patents, Washington, D.C. 20591

St. St. Andrew
Printed Name

St. St. Andrew
Signature

7. ☒ Preliminarily, please amend the specification by inserting before the first line the following:

--This application is a ☐ continuation ☒ division of application Serial No. 08/535,681, filed September 28, 1995, which is hereby incorporated by reference.--

8. ☐ Transfer the drawings from the prior application to this application and abandon said prior application as of the filing date accorded this application. A duplicate copy of this sheet is enclosed for filing in the prior application file. (May only be used if signed by person authorized by § 1.138 and before payment of base issue fee.)

8a. ☒ New formal drawings are enclosed.

9. ☐ Priority of application Serial No. _____ filed on _____ in _____ (country) is claimed under 35 USC 119.

9a. ☐ The certified copy has been filed in prior application Serial No. _____ filed _____

10. ☒ The prior application is assigned of record to Amgen Inc.

11. ☒ A preliminary amendment is enclosed.

12. ☒ Also enclosed Attorney's Statement, Information Disclosure Statement, Modified Form 1449

13. ☐ Other: _____

14. ☒ The power of attorney in the prior application is to:
Ron K. Levy, Registration No.: 31,539; Steven M. Odre, Registration No.: 29,094;
Daniel R. Curry, Registration No.: 32,727


a. ☒ The power appears in the original papers in the prior application.

b. ☐ Since the power does not appear in the original papers, a copy of the power in the prior application is enclosed.

c. ☒ Address all future communications to
Daniel R. Curry
at the address below.

Signator: ☐ Assignee of complete interest

☒ Attorney or agent of record



Daniel R. Curry
Attorney for Applicant
Registration No. 32,727
Phone: (805) 447-8102
Date: October 13, 2000

Please send all future correspondence to:

U. S. Patent Operations/ DRC
Dept. 4300, M/S 27-4-A
AMGEN, INC.
One Amgen Center Drive
Thousand Oaks, California 91320-1799

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant(s): Shaw-Fen Sylvia Hu

Serial No.: To Be Assigned

Group Art Unit No.: Not Yet Assigned

Filed: October 13, 2000

Examiner: Not Yet Assigned

For: TRUNCATED GLIAL CELL LINE-DERIVED
NEUROTROPHIC FACTOR

Docket No.: A-357B

PRELIMINARY AMENDMENTAssistant Commissioner for Patents
Washington, D. C. 20231

Sir:

Please amend the referenced application as follows:

In the Specification

Page 1, line 2, insert -- This application is a divisional of application Serial No. 08/535,681, filed September 28, 1995 now pending, which is hereby incorporated by reference. --

In the Claims

Please cancel Claims 1-29 and 31-44.

Please amend the following claims:

30. (Amended) A method for affecting the survival or function of neurons [of treating Parkinson's Disease] comprising administering [to a patient the] a pharmaceutical composition [of Claim 27] comprising:

EXPRESS MAIL CERTIFICATE"Express Mail" mail labeling number: EL360889268USDate of Deposit: October 13, 2000

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to Box Patent Application, Assistant Commissioner for Patents, Washington, DC 20231

Sh. Sh. Anderson
Printed Name

Sh. Sh. Anderson
Signature

(a) a truncated glial cell line-derived neurotrophic factor (GDNF) protein product consisting of an amino acid sequence

X-(Cys⁴¹-Cys¹³³)-Y

wherein

(Cys⁴¹-Cys¹³³) consists of Cys⁴¹ through Cys¹³³ of SEQ ID NO:2;

Y represents the carboxy terminal group of Cys¹³³, a carboxy-terminus amino acid residue of Ile¹³⁴, or a substituted amino acid residue, and

X represents a methionylated or nonmethionylated amine group of Cys⁴¹ or amino-terminus amino acid residue(s) selected from the group:

G
RG
NRG
KNRG (SEQ ID NO:3)
GKNRG (SEQ ID NO:4)
RGKNRG (SEQ ID NO:5)
QRGKNRG (SEQ ID NO:6)
GORGKNRG (SEQ ID NO:7)
RGORGKNRG (SEQ ID NO:8)
RRGORGKNRG (SEQ ID NO:9)
G RRGORGKNRG (SEQ ID NO:10)
KG RRGORGKNRG (SEQ ID NO:11)
GKG RRGORGKNRG (SEQ ID NO:12)
RGKG RRGORGKNRG (SEQ ID NO:13)
SRGKG RRGORGKNRG (SEQ ID NO:14)
NSRGKG RRGORGKNRG (SEQ ID NO:15)
ENSRGKG RRGORGKNRG (SEQ ID NO:16)
PENSRGKG RRGORGKNRG (SEQ ID NO:17)
NPENSRGKG RRGORGKNRG (SEQ ID NO:18)
ANPENSRGKG RRGORGKNRG (SEQ ID NO:19)
A ANPENSRGKG RRGORGKNRG (SEQ ID NO:20)
AA ANPENSRGKG RRGORGKNRG (SEQ ID NO:21)
AAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:22)
OAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:23)
ROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:24)

NROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:25)

RNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:26)

ERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:27)

RERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:28)

RRERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:29)

P RRERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:30)

LP RRERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:31)

VL P RRERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:32)

AVLP RRERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:33)

MAVLP RRERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:34)

OMAVLP RRERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:35)

KOMAVLP RRERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:36)

DKOMAVLP RRERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:37) and

PDKOMAVLP RRERNROAAA ANPENSRGKG RRGORGKNRG (SEQ ID NO:38)


or a substitution or deletion variant of X, wherein said variant is in excess of 70% identical to an amino acid sequence of X as set forth above when four gaps in a length of 100 amino acids may be introduced to assist in that alignment, and

(b) a pharmaceutically acceptable vehicle.

Please add the following claims:

-- 45. **(Newly added)** A method for affecting the survival or function of neurons comprising administering a pharmaceutical composition comprising:

(a) a truncated glial cell line-derived neurotrophic factor (GDNF) protein product consisting of an amino acid sequence



wherein

(Cys⁴¹-Cys¹³³) consists of Cys⁴¹ through Cys¹³³ of SEQ ID NO:2;

Y represents the carboxy terminal group of Cys¹³³, a carboxy-terminus amino acid residue of Ile¹³⁴, or a substituted amino acid residue, and

X represents a methionylated or nonmethionylated amine group of Cys⁴¹ or amino-terminus amino acid residue(s) selected from the group:

G
 RG
 NRG
 KNRG (SEQ ID NO:3)
 GKNRG (SEQ ID NO:4)
 RGKNRG (SEQ ID NO:5)
 QRGKNRG (SEQ ID NO:6)
 GQRGKNRG (SEQ ID NO:7)
 RGQRGKNRG (SEQ ID NO:8)
 RRGQRGKNRG (SEQ ID NO:9)
 G RRGQRGKNRG (SEQ ID NO:10)
 KG RRGQRGKNRG (SEQ ID NO:11)
 GKG RRGQRGKNRG (SEQ ID NO:12)
 RGKG RRGQRGKNRG (SEQ ID NO:13)
 SRGKG RRGQRGKNRG (SEQ ID NO:14)
 NSRGKG RRGQRGKNRG (SEQ ID NO:15)
 ENSRGKG RRGQRGKNRG (SEQ ID NO:16)
 PENSRGKG RRGQRGKNRG (SEQ ID NO:17)
 NPENSRGKG RRGQRGKNRG (SEQ ID NO:18)
 ANPENSRGKG RRGQRGKNRG (SEQ ID NO:19)
 A ANPENSRGKG RRGQRGKNRG (SEQ ID NO:20)
 AA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:21)
 AAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:22)
 QAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:23)
 RQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:24)
 NRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:25)
 RNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:26)
 ERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:27)
 RERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:28)
 RRERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:29)
 P RRERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:30)
 LP RRERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:31)
 VLP RRERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:32)
 AVL P RRERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:33)
 MAVLP RRERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:34)
 QMAVLP RRERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:35)
 KQMAVLP RRERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:36)
 DKQMAVLP RRERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:37) and

PDKQMAVL P RRERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:38); and

(b) a pharmaceutically acceptable vehicle.

46. A method according to Claim 30 or 45, wherein X is selected from the group consisting of SEQ ID NO: 3, 7, 8, 14, 17 and 18.

47. A method according to Claim 30 or 45, wherein X is G, RG or NRG.

48. A method according to Claim 30 or 45, wherein said GDNF protein product has the amino acid sequence of SEQ ID NO:42.

49. A method according to Claim 30 or 45, wherein said GDNF protein product has the amino acid sequence of SEQ ID NO:44.

50. A method according to Claim 30 or 45, wherein said GDNF protein product has the amino acid sequence of SEQ ID NO:46. --

Respectfully submitted,



Daniel R. Curry
Attorney for Applicant
Registration No.: 32,727
Phone: (805) 447-8102
Date: October 13, 2000

Please send all future correspondence to:

U.S. Patent Operations/ DRC
Dept. 4300, M/S 27-4-A
AMGEN INC.
One Amgen Center Drive
Thousand Oaks, California 91320-1799

TRUNCATED GLIAL CELL LINE-DERIVED NEUROTROPHIC FACTOR

Field of the Invention

In general, the present invention relates to proteins, referred to herein as glial cell line-derived neurotrophic factors (also referred to as glial derived neurotrophic factor or GDNF), that are characterized by the ability to promote dopamine uptake by dopaminergic neurons and support the survival of the neurons that die in Parkinson's Disease. The present invention more specifically relates to a novel truncated GDNF proteins.

Background of the Invention

Neurotrophic factors are proteins, found in the nervous system or in non-nerve tissues innervated by the nervous system, whose function is to promote the survival and maintain the phenotypic differentiation of nerve and/or glial cells (Varon et al., *Ann. Rev. Neuroscience* 1:327, 1979; Thoenen et al., *Science* 229:238, 1985). Because of this physiological role, neurotrophic factors are useful in treating the degeneration of nerve cells and the loss of differentiated function that occurs in a variety of neurodegenerative diseases.

In order for a particular neurotrophic factor to be potentially useful in treating nerve damage, the class or classes of damaged nerve cells must be responsive to the factor. Different neurotrophic factors typically affect distinctly different classes of nerve cells. Therefore, it is advantageous to have on hand a variety of different neurotrophic factors to treat each of the classes of damaged neurons that may occur with different forms of disease or injury.

Neurotrophic factors can protect responsive neurons against a variety of unrelated insults. For example, nerve growth factor (NGF) will rescue a significant portion of sensory neurons from death caused by cutting their axonal processes (Rich et al., *J. Neurocytol* 16:261, 1987; Otto et al., *J. Neurosci.* 83:156, 1987), from ontogenetic death during embryonic development (Hamburger et al., *J. Neurosci.* 4:767, 1984), and from damage caused by the administration of taxol or cisplatin (Apfel et al., *Ann. Neurol.* 29: 87, 1991). This apparent generality of protection has led to the concept that if a neurotrophic factor protects responsive neurons against experimental damage, it may be useful in treating diseases that involve damage to those neurons in patients, even though the etiology may be unknown.

Express Mail [®] mail labeling number: TB 530289575 US		Date of Deposit: September 28, 1995
<small>I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. 1.10 on the date indicated above and is addressed to Box Patent Application, Assistant Commissioner of Patents, Washington, D. C. 20231. (Senders Signature Below)</small>		
(typed name) Honda G. Spahr		<i>Honda G. Spahr</i>

A given neurotrophic factor, in addition to having the correct neuronal specificity, must be available in sufficient quantity to be used as a pharmaceutical treatment. Since neurotrophic factors are typically present in small amounts in tissues (e.g., Hofer and Barde *Nature* 331:261, 1988; Lin et al., *Science* 246:1023, 1989), it would be inconvenient to prepare pharmaceutical quantities of neurotrophic factors directly from animal tissues. As an alternative, it is desirable to use a recombinant expression system to produce the desired protein.

Lin et al. previously described a method for screening biological samples for neurotrophic activity on the embryonic precursors of the substantia nigra dopaminergic neurons (see U.S. Patent Application No. 08/182,183 filed May 23, 1994 and its parent applications; PCT/US92/07888 filed September 17, 1992 (WO 93/06116); and European Patent Application No. 92921022.7 (Publication No. EP 610 254); the disclosures of which are hereby incorporated by reference). This bioassay is useful in identifying neurotrophic factors which may be used in treating Parkinson's disease (Friedman et al., *Neuro. Sci. Lett.* 79:65-72, 1987) as the disease is characterized by the degeneration of dopaminergic neurons in the midbrain that innervate the striatum.

Lin et al. further described the characterization of a new neurotrophic factor that was purified from one such source, the conditioned culture medium from a glioblastoma cell line, B49 (Schubert et al., *Nature* 249:224-27, 1974). The conditioned medium from this cell line was previously reported to contain dopaminergic neurotrophic activity (Bohn et al., *Soc. Neurosci. Abs.* 15:277, 1989). Prior to the disclosure of Lin et al., glial cell line-derived neurotrophic factor (GDNF) had not been identified as a discrete biologically active substance or isolated as a substantially pure protein. In addition, Lin et al. described processes for cloning human genes encoding GDNF, the nucleic acid sequence of the human genes that encode GDNF and the amino acid sequences of the GDNF protein. The GDNF gene was subcloned into an expression vector, and the vector was used to express biologically active GDNF. The GDNF protein is a homodimer composed of two 134 amino acid, 22 kDa, subunits joined by disulfide bond. The description further included the use of GDNF for preventing and treating nerve damage and nerve related diseases such as Parkinson's disease.

GDNF therapy is helpful in the treatment of nerve damage caused by conditions that compromise the survival and/or proper function of one or more types of nerve cells. Such nerve damage may occur from a wide variety of

different causes. Nerve damage may occur to one or more types of nerve cells by: (1) physical injury, which causes the degeneration of the axonal processes and/or nerve cell bodies near the site of injury; (2) temporary or permanent cessation of blood flow to parts of the nervous system, as in stroke;

- 5 (3) intentional or accidental exposure to neurotoxins, such as chemotherapeutic agents (e.g., cisplatin) for the treatment of cancer or dideoxycytidine (ddC) for the treatment of AIDS; (4) chronic metabolic diseases, such as diabetes or renal dysfunction; or (5) neurodegenerative diseases such as Parkinson's disease, Alzheimer's disease, and Amyotrophic Lateral Sclerosis (ALS), which
10 result from the degeneration of specific neuronal populations.

GDNF therapy could be particularly helpful in the treatment of neurodegenerative conditions involving the degeneration of the dopaminergic neurons of the substantia nigra, such as Parkinson's disease. The only current treatments for Parkinson's disease are palliative, aiming at increasing dopamine
15 levels in the striatum. The expected impact of GDNF therapy is not simply to produce an increase in the dopaminergic neurotransmission at the dopaminergic nerve terminals in the striatum (which will result in a relief of the symptoms), but also to slow down, or even stop, the progression of the degenerative processes and to repair the damaged nigrostriatal pathway and restore its function. GDNF
20 may also be used in treating other forms of damage to or improper function of dopaminergic nerve cells in human patients. Such damage or malfunction may occur in schizophrenia and other forms of psychosis. The only current treatments for such conditions are symptomatic and require drugs which act upon dopamine receptors or dopamine uptake sites, consistent with the view that
25 the improper functioning of the dopaminergic neurons which innervate these receptor-bearing neuronal populations may be involved in the disease process.

SUMMARY OF THE INVENTION

30

In one aspect, the present invention provides novel truncated glial cell line-derived neurotrophic factor (GDNF) protein products. In one embodiment, truncated GDNF proteins are produced by recombinant genetic engineering techniques. In an alternative embodiment, the truncated GDNF proteins are
35 synthesized by chemical techniques, or a combination of the recombinant and chemical techniques.

The truncated GDNF protein products of the present invention include the proteins represented by the amino acid sequence X-[Cys⁴¹-Cys¹³³]-Y. The amino acid residue numbering scheme of Figure 1 (SEQ ID NO:2) is used to facilitate comparison to the mature GDNF protein. [Cys⁴¹-Cys¹³³] represents the amino acid sequence of Cys⁴¹ through Cys¹³³ as depicted in Figure 1 (SEQ ID NO:2). Y represents the carboxy terminal group of Cys¹³³ or a carboxy-terminus amino acid residue of Ile¹³⁴. X represents a methionylated or nonmethionylated amine group of Cys⁴¹ or amino-terminus amino acid residue(s) selected from the group:

10

	G
	RG
	NRG
	KNRG (SEQ ID NO:3)
	GKNRG (SEQ ID NO:4)
	RGKNRG (SEQ ID NO:5)
	QRGKNRG (SEQ ID NO:6)
	GQRGKNRG (SEQ ID NO:7)
	RGQRGKNRG (SEQ ID NO:8)
	RRGQRGKNRG (SEQ ID NO:9)
G	RRGQRGKNRG (SEQ ID NO:10)
KG	RRGQRGKNRG (SEQ ID NO:11)
GKG	RRGQRGKNRG (SEQ ID NO:12)
RGKG	RRGQRGKNRG (SEQ ID NO:13)
SRGKG	RRGQRGKNRG (SEQ ID NO:14)
NSRGKG	RRGQRGKNRG (SEQ ID NO:15)
ENSRGKG	RRGQRGKNRG (SEQ ID NO:16)
PENSRGKG	RRGQRGKNRG (SEQ ID NO:17)
NPENSRGKG	RRGQRGKNRG (SEQ ID NO:18)
ANPENSRGKG	RRGQRGKNRG (SEQ ID NO:19)
A	ANPENSRGKG RRGQRGKNRG (SEQ ID NO:20)
AA	ANPENSRGKG RRGQRGKNRG (SEQ ID NO:21)
AAA	ANPENSRGKG RRGQRGKNRG (SEQ ID NO:22)
QAAA	ANPENSRGKG RRGQRGKNRG (SEQ ID NO:23)
RQAAA	ANPENSRGKG RRGQRGKNRG (SEQ ID NO:24)
NRQAAA	ANPENSRGKG RRGQRGKNRG (SEQ ID NO:25)

	RNRQAAA	ANPENSRRGKG	RRGQRGKNRG	(SEQ ID NO:26)
	ERNRQAAA	ANPENSRRGKG	RRGQRGKNRG	(SEQ ID NO:27)
	RERNRQAAA	ANPENSRRGKG	RRGQRGKNRG	(SEQ ID NO:28)
	RRERNRQAAA	ANPENSRRGKG	RRGQRGKNRG	(SEQ ID NO:29)
P	RRERNRQAAA	ANPENSRRGKG	RRGQRGKNRG	(SEQ ID NO:30)
LP	RRERNRQAAA	ANPENSRRGKG	RRGQRGKNRG	(SEQ ID NO:31)
VLP	RRERNRQAAA	ANPENSRRGKG	RRGQRGKNRG	(SEQ ID NO:32)
AVLP	RRERNRQAAA	ANPENSRRGKG	RRGQRGKNRG	(SEQ ID NO:33)
MAVLP	RRERNRQAAA	ANPENSRRGKG	RRGQRGKNRG	(SEQ ID NO:34)
QMAVLP	RRERNRQAAA	ANPENSRRGKG	RRGQRGKNRG	(SEQ ID NO:35)
KQMAVLP	RRERNRQAAA	ANPENSRRGKG	RRGQRGKNRG	(SEQ ID NO:36)
DKQMAVLP	RRERNRQAAA	ANPENSRRGKG	RRGQRGKNRG	(SEQ ID NO:37)
PDKQMAVLP	RRERNRQAAA	ANPENSRRGKG	RRGQRGKNRG	(SEQ ID NO:38)

It is contemplated that such truncated GDNF protein products would include truncated GDNF protein having the amino acid sequence as represented by X-[Cys⁴¹-Cys¹³³]-Y and variants and derivatives thereof. Thus, the truncated

5 GDNF protein products of the present invention also include addition, substitution and internal deletion variants and derivatives of the amino acid sequences represented by X-[Cys⁴¹-Cys¹³³]-Y. The truncated GDNF protein products further include methionylated or nonmethionylated forms as well as glycosylated or non-glycosylated forms of truncated GDNF protein.

10 Exemplary truncated GDNF proteins of the present invention include [Arg¹⁶-Ile¹³⁴], [Asn²²-Ile¹³⁴], [Pro²³-Ile¹³⁴], [Ser²⁶-Ile¹³⁴], [Arg³²-Ile¹³⁴], [Gly³³-Ile¹³⁴], [Lys³⁷-Ile¹³⁴] and [Asn³⁸-Ile¹³⁴] truncated GDNF proteins, either methionylated or nonmethionylated, and variants and derivatives thereof. Presently preferred truncated GDNF proteins

15 of the present invention include [Lys³⁷-Ile¹³⁴] and [Asn³⁸-Ile¹³⁴] truncated GDNF proteins, either methionylated or nonmethionylated, and variants and derivatives thereof. Exemplary substitution variants are the [Asn²²ΔSer²²-Ile¹³⁴] and [Pro²³-Lys³⁷ΔAsn³⁷-Ile¹³⁴] truncated GDNF proteins. An exemplary addition variant is the Ser-[Pro²³-Ile¹³⁴] truncated GDNF protein.

20 In another aspect of the present invention, the truncated GDNF proteins may be made in glycosylated or non-glycosylated forms. Derivatives of truncated GDNF protein typically involve attaching the truncated GDNF protein to a water soluble polymer. For example, the truncated GDNF protein may be conjugated to

one or more polyethylene glycol molecules to decrease the precipitation of the truncated GDNF protein product in an aqueous environment.

Yet another aspect of the present invention includes the various polynucleotides encoding truncated GDNF proteins. These nucleic acid sequences are generally used in the expression of truncated GDNF in a eukaryotic or prokaryotic host cell, wherein the expression product or a derivative thereof is characterized by the ability to increase dopamine uptake by dopaminergic cells. The polynucleotides may also be used in cell therapy or gene therapy applications. Suitable nucleic acid sequences include those specifically depicted in the Figures as well as additional degenerate sequences and naturally occurring allelic variations.

A further aspect of the present invention involves vectors containing the polynucleotides encoding truncated GDNF proteins operatively linked to amplification and/or expression control sequences. Both prokaryotic and eukaryotic host cells may be stably transformed or transfected with such vectors to express the truncated glial derived neurotrophic factor. The present invention further includes the recombinant production of a truncated GDNF protein wherein such transformed or transfected host cells are grown in a suitable nutrient medium, and the truncated GDNF expressed by the cells is, optionally, isolated from the host cells and/or the nutrient medium. The present invention further includes the use of polynucleotides encoding truncated GDNF and vectors containing such polynucleotides in gene therapy or cell therapy.

In another aspect, the present invention involves a recombinantly produced GDNF composition containing a mixture of a mature GDNF protein and one or more truncated GDNF proteins derived therefrom, wherein the mature GDNF protein has a molecular weight of approximately 44 kDa, and wherein the truncated GDNF protein has a molecular weight of approximately 36 to 40 kDa. The GDNF composition may contain at least two truncated GDNF species wherein a first species has a molecular weight of approximately 36 kDa and a second species has a molecular weight of approximately 40 kDa. The truncated GDNF species having a molecular weight of approximately 40 kDa is a heterodimer of a GDNF monomer having a molecular weight of approximately 22 kDa and a truncated GDNF monomer having a molecular weight of approximately 18 kDa. It is also contemplated that one or more of the truncated GDNF species may be isolated from such a mixture for therapeutic use.

Another aspect of the present invention includes pharmaceutical compositions containing truncated GDNF protein product. Typically, the truncated GDNF protein product is formulated in association with a pharmaceutically acceptable vehicle. A variety of other formulation materials may be used to facilitate manufacture, storage, handling, delivery and/or efficacy. In another aspect of the present invention, truncated GDNF protein products increase dopamine uptake and survival of dopaminergic neurons. Thus, the truncated GDNF protein products are particularly suitable for the treatment of damage to the nervous system caused by injury or disease, such as Parkinson's Disease.

Additional aspects and advantages of the invention will be apparent to those skilled in the art upon consideration of the following description, which details the practice of the present invention.

BRIEF DESCRIPTION OF THE FIGURES

Numerous features and advantages of the present invention will become apparent upon review of the figures, wherein:

Figure 1 depicts a nucleic acid sequence (SEQ ID NO:1) encoding mature human glial cell line-derived neurotrophic factor (hGDNF). Also depicted is the amino acid sequence (SEQ ID NO:2) of the mature human GDNF protein.

Figure 2 depicts a diagram of a plasmid construction made for the expression of recombinant truncated GDNF proteins.

Figure 3 depicts a restriction map of an alternative nucleic acid sequence (SEQ ID NO:39) encoding GDNF and truncated GDNF polynucleotides.

Figure 4 depicts a restriction map of yet another nucleic acid sequence (SEQ ID NO:40) encoding GDNF and truncated GDNF polynucleotides.

Figure 5 depicts a nucleic acid sequence (SEQ ID NO:41) encoding [Pro²³-Lys³⁷ΔAsn³⁷-Ile¹³⁴] truncated GDNF protein substitution variant

(SEQ ID NO:42). This protein may also be described as a Met-Ser-[Pro²³-Lys³⁷ Δ Asn³⁷-Ile¹³⁴] truncated GDNF protein addition/substitution variant.

Figure 6 depicts a nucleic acid sequence (SEQ ID NO:43) encoding an
5 [Arg³²-Ile¹³⁴] truncated GDNF protein (SEQ ID NO:44).

Figure 7 depicts a nucleic acid sequence (SEQ ID NO:45) encoding a
[Gly³³-Ile¹³⁴] truncated GDNF protein (SEQ ID NO:46).

10 Figure 8 depicts the amino acid sequence of mature hGDNF (SEQ ID NO:47) in comparison to several exemplary truncated GDNF proteins: Met-[Arg³²-Ile¹³⁴] (SEQ ID NO:48), Met-[Gly³³-Ile¹³⁴] (SEQ ID NO:49) and Met-Ser-[Pro²³-Lys³⁷ Δ Asn³⁷-Ile¹³⁴] (SEQ ID NO:50).

15 DETAILED DESCRIPTION OF THE INVENTION

Human glial cell line-derived neurotrophic factor (hGDNF) is synthesized as a precursor that is processed and secreted as a mature protein of
20 134 amino acids. It was previously determined that mature human GDNF has the amino acid sequence depicted in Figure 1 (SEQ ID NO:2).

The present invention is based on the unexpected discovery that the mature GDNF protein may be reduced in size (also referred to herein as a "clipped" or "truncated" protein or truncated GDNF protein) yet retain its
25 biological activity. The clipped protein was first discovered during the recombinant production of GDNF in Chinese hamster ovary (CHO) cells. In brief, the recombinant human GDNF (rhGDNF) was prepared as follows. A nucleic acid sequence encoding the entire open reading frame of the mature human GDNF protein was cloned into an expression plasmid. The nucleic acid sequence was
30 confirmed to be correct (by DNA sequencing as the equivalent of the hGDNF sequence in GeneBank) and was translated to an amino acid sequence identical to the published sequence for mature human GDNF (Lin et al., *Science* 260, 1130-1132, 1993). The plasmid DNA was linearized and transfected into dihydrofolate reductase-deficient CHO cells (CHO⁻ cells) using the calcium
35 phosphate precipitation method. Transfected cells were cultured in a selective

medium, and those colonies that survived the selection process were chosen for individual analysis of hGDNF expression.

Serum-free conditioned media from the individual clones were collected and subjected to Western blot analysis using antisera specific for hGDNF. The antisera involved rabbit polyclonal antibodies elicited from rabbits immunized with recombinant hGDNF expressed in *Escherichia coli*. Under reducing conditions, the hGDNF that was present in these samples was resolved into two major bands having apparent molecular weights of approximately 22 kDa and 18 kDa. Each band consisted of a closely spaced doublet of approximately 22 + 22.5 kDa and 18 + 17.5 kDa, respectively (for simplicity, these doublets will be referred as the 22 kDa and 18 kDa bands or species).

GDNF had previously been reported to exist as a disulfide-bonded homodimer composed of two identical subunits of the mature GDNF protein having a molecular weight of approximately 20 to 22 kDa. When GDNF was analyzed under nonreducing conditions, it was reported that a broad band of 32 to 42 kDa (Lin et al., *Science* 260, 1130-1132, 1993) or 33 to 45 kDa (Lin et al., *J. Neurochem.* 63(2), 758-768, 1994) had been identified. The existence of the range was interpreted as being due to the heterogeneity of glycosylation on the mature monomers and was further substantiated with de-glycosylation experiments.

While the present 22 kDa band corresponds to the mature GDNF protein reported in the literature, the 18 kDa band has not previously been reported. The relative amounts of the 22 kDa and the 18 kDa protein varied in samples collected from individual clones. In addition, it was found that multiple harvests from the same clone showed a variable ratio of the two bands. Moreover, it was found that storage of the CHO-expressed GDNF protein frequently led to an increase in the presence of the 18 kDa band with a concurrent decrease of the 22 kDa band.

When conditioned medium from the transformed CHO⁻ cells was analyzed under nonreducing conditions by Western blots, three well-resolved bands with apparent molecular weights of 36, 40 and 44 kDa were observed. This finding was also in contrast to previous reports. The relative intensity of these bands was variable, but they correlated well with the ratio of the 22 and 18 kDa monomer bands present in each of the samples. Upon further analysis with monoclonal antisera, it was determined that the three bands in the nonreducing gel corresponded to three possible dimers composed of the two monomers. The

largest 44 kDa protein is a dimer of two 22 kDa mature GDNF proteins as previously reported. The intermediate 40 kDa protein consists of a dimer in which one mature protein has been reduced in molecular weight to an 18 kDa form. The smallest 36 kDa dimer appears to contain two 18 kDa proteins, i.e.,

both 22 kDa forms have been reduced in molecular weight. This data demonstrated for the first time not only the presence of a novel form of GDNF monomer but also the presence of the clipped GDNF protein in the dimeric configuration. It was also found that, when stored, the monomer composition of the samples shifted towards that of the clipped form and the corresponding dimer species, i.e., the amount of the 36 kDa protein was seen to increase.

Studies were then performed to identify which part of the protein was being eliminated or changed to cause of the reduction in molecular weight in comparison to that of the previously reported mature GDNF protein. It was first determined that the reduction in molecular weight was not due to changes in glycosylation.

GDNF contains two potential N-linked glycosylation sites and has been reported to be glycosylated. The clipped protein, however, is not simply the nonglycosylated or underglycosylated form of mature GDNF. This was demonstrated in deglycosylation experiments wherein samples were treated with N-glycanase, O-glycanase and neuraminidase. On reducing gels, the 18 kDa protein was reduced to a 13.5 kDa band by N-glycanase digestion indicating the presence of an equivalency of 4.5 kDa of N-linked sugar. Treatment with neuraminidase and O-glycanase caused the 18 kDa band to shift slightly to 17 kDa. This indicated the presence of O-linked sugars on the protein. The mature 22 kDa band has been reported to be glycosylated and was also reduced to 18 kDa (i.e., also by 4.5 kDa) by N-glycanase. This was further confirmed through the use of a monoclonal antibody which is specific for the 22 kDa band on the gel. The glycanase digestion pattern of the nonreduced dimer was more complicated, but was interpretable and consistent with the initial assignment of the three forms.

As a result, the 4.5 kDa reduction in molecular weight of the protein was then viewed as resulting from the deletion of approximately 30-35 amino acid residues rather than from changes in glycosylation. The deletion was expected to most likely occur at the amino-terminus of the mature GDNF protein for the following reasons. Mature GDNF contains a total of seven cystines. If the deletion were from the carboxyl terminus, 2 to 4 of the seven cystines would be lost, and

this would likely result in an inactive protein. However, when a test sample consisting of predominantly the clipped form was subjected to a bioassay to measure its dopaminergic neuron neurotrophic activity, the sample demonstrated comparable activity to a sample which contained proportionally more of the mature form of GDNF.

The site of cleavage was then determined via amino acid sequence analysis of the purified protein. Samples were sequenced, according to manufacturer's instructions, using an Applied Biosystems 494A protein sequencer for ten cycles. While amino acid sequence analysis techniques and procedures are well known to those skilled in the art, further descriptions of the sequencing of proteins are provided in Fausset et al., *Electrophoresis* 12:22-27, 1991 and United States Patent Application No. 576.316 filed August 24, 1990 (European Patent Application No. 90310899, Publication No. EP 423 980, filed October 4, 1990, entitled "Stem Cell Factor") the disclosures of which are hereby incorporated by reference. Upon analysis, it was determined that the amino terminus of the clipped protein was "RGQRGK" or Arg-Gly-Gln-Arg-Gly-Lys. Therefore, the first 31 amino acids had been removed from the mature protein in the conditioned media. The remaining amino acid sequence of the clipped protein, beginning with amino acid Arg³², was otherwise consistent with that of the mature GDNF amino acid sequence depicted in Figure 1 (SEQ ID NO:2).

The [Arg³²-Ile¹³⁴] truncated GDNF protein was found to be active, on a qualitative basis, in a dopaminergic neuron assay. The dopaminergic neurotrophic activity assay is used to identify neurotrophic factors that may be beneficial in treating Parkinson's disease. The assay is based on a previously described assay (Friedman et al., *Neuro. Sci. Lett.* 79:65-72, 1987, the disclosure of which is hereby incorporated by reference) and may include modifications as described in Lin et al. (see U.S. Patent Application No. 08/182,183 filed May 23, 1994 and its parent applications; PCT/US92/07888 filed September 17, 1992 (WO 93/06116); and European Patent Application No. 92921022.7 (Publication No. EP 610 254)). A detailed description of the assay is provided in Example 5, below.

A subsequent purification procedure, followed by amino acid sequencing, led to the discovery of another protein from which the first 36 amino acid residues had been removed from the N-terminus of mature GDNF: a [Lys³⁷-Ile¹³⁴] truncated GDNF protein, with an N-terminal sequence of KNRG(C)VL--. Again, the remaining amino acid residues of the clipped protein were otherwise

consistent with those of the mature human GDNF amino acid sequence. The [Lys³⁷-Ile¹³⁴] truncated GDNF protein was also analyzed in the dopamine uptake bioassay. This truncated GDNF protein was found to be active with an ED₅₀ of about 50 pg/ml, similar to that of purified recombinant *E. coli*-expressed, mature GDNF.

It was further discovered that bacterially expressed mature GDNF could be changed to a truncated form. Mature GDNF, expressed in transformed *E. coli* (as described in Lin et al., U.S. Patent Application No. 08/182,183, *supra*), was incubated with CHO-derived conditioned media. Recombinant *E. coli* GDNF has an apparent molecular weight of 17 kDa on reducing gel. When the material was mixed with CHO cell conditioned media and incubated for five days at 4°C, the protein was clipped completely to 12.5 kDa. This cleavage was less complete with one hour or 24 hours of incubation, suggesting a time-dependent process under such conditions. It was also found that simply incubating recombinant *E. coli* GDNF overnight with media containing 0.1% fetal bovine serum did not generate the clipped form. Thus, the presence of live cells in the culture seems to be necessary for the clipping process to occur. It is possible, therefore, that the clipping event may also occur *in vivo* within certain tissues.

In addition, it was found that derivatives of mature *E. coli*-expressed hGDNF, such as pegylated GDNF (also described in Lin et al., U.S. Patent Application No. 08/182,183, *supra*) may be processed to a truncated form in the presence of CHO-derived conditioned media. Mature GDNF may be pegylated at the amino terminus in order to enhance its clearance time in circulation. Pegylation increases the size of the protein, and the modified mature GDNF migrates at about 45 kDa under reduced conditions. As with the non-pegylated mature form, the incubation of pegylated *E. coli* GDNF with CHO cell (untransfected) conditioned media generated a 12.5 kDa band. In both cases, the 12.5 kDa species was present as a disulfide-bonded dimer as shown on non-reducing gels. The generation of this clipped form from the N-terminally pegylated mature protein further demonstrated that the clipping event occurred at the N-terminus of the protein since the pegylated residue was lost during the clipping process.

Based on these findings and because the clipping event may also occur *in vivo*, a truncated form of the GDNF protein may be the ultimate naturally processed form of hGDNF under physiological conditions. Therefore, it was considered advantageous to produce a truncated GDNF protein, or derivative

thereof, for therapeutic use. For example, a directly expressed or synthesized truncated GDNF protein, such as the [Arg³²-Ile¹³⁴] truncated GDNF protein, would be expected to be resistant to the above-described proteolytic activity. Moreover, if it was desired to produce a truncated GDNF derivative, such as a

5 pegylated [Arg³²-Ile¹³⁴] truncated GDNF protein, the resulting derivative would be expected to have the advantage of not being susceptible to the specific clipping which was observed with the mature GDNF derivative.


Additional advantages can also be expected of truncated GDNF protein products. First, the pI of a truncated protein, such as the [Arg³²-Ile¹³⁴]

10 truncated GDNF protein, will be reduced from about 10 to about 8.0-8.5. This makes the protein significantly less basic which could in turn provide beneficial effects including better receptor binding and decreased cytotoxicity at the site of administration, such as an intrathecal injection site. Second, within the first 26 amino acids of the mature GDNF amino acid sequence are two deamidation sites:

15 Arg-Asn-Arg (amino acids 14-16) and Glu-Asn-Ser (amino acids 24-26). The absence of one or both of these sites in a truncated GDNF protein is expected to increase the stability of the protein.

Truncated GDNF Protein Products

20 In a basic embodiment, the truncated GDNF proteins of the present invention may be represented by the following amino acid sequence wherein the amino acid residue numbering scheme of Figure 1 is used to facilitate comparison to the mature GDNF protein:



25 wherein

[Cys⁴¹-Cys¹³³] represents the amino acid sequence of Cys⁴¹ through Cys¹³³ as depicted in Figure 1 (SEQ ID NO:2);

Y represents the carboxy terminal group of Cys¹³³ or a carboxy-terminus amino acid residue of Ile¹³⁴; and

30 X represents a methionylated or nonmethionylated amine group of Cys⁴¹ or amino-terminus amino acid residue(s) selected from the group:

G
RG
NRG
KNRG (SEQ ID NO:3)

GKNRG (SEQ ID NO:4)
 RGKNRG (SEQ ID NO:5)
 QRGKNRG (SEQ ID NO:6)
 GQRGKNRG (SEQ ID NO:7)
 RGQRGKNRG (SEQ ID NO:8)
 RRQRGKNRG (SEQ ID NO:9)
 G RRQRGKNRG (SEQ ID NO:10)
 KG RRQRGKNRG (SEQ ID NO:11)
 GKG RRQRGKNRG (SEQ ID NO:12)
 RGKG RRQRGKNRG (SEQ ID NO:13)
 SRKG RRQRGKNRG (SEQ ID NO:14)
 NSRKG RRQRGKNRG (SEQ ID NO:15)
 ENSRKG RRQRGKNRG (SEQ ID NO:16)
 PENSRKG RRQRGKNRG (SEQ ID NO:17)
 NPENSRKG RRQRGKNRG (SEQ ID NO:18)
 ANPENSRKG RRQRGKNRG (SEQ ID NO:19)
 A ANPENSRKG RRQRGKNRG (SEQ ID NO:20)
 AA ANPENSRKG RRQRGKNRG (SEQ ID NO:21)
 AAA ANPENSRKG RRQRGKNRG (SEQ ID NO:22)
 QAAA ANPENSRKG RRQRGKNRG (SEQ ID NO:23)
 RQAAA ANPENSRKG RRQRGKNRG (SEQ ID NO:24)
 NRQAAA ANPENSRKG RRQRGKNRG (SEQ ID NO:25)
 RNRQAAA ANPENSRKG RRQRGKNRG (SEQ ID NO:26)
 ERNRQAAA ANPENSRKG RRQRGKNRG (SEQ ID NO:27)
 RERNRQAAA ANPENSRKG RRQRGKNRG (SEQ ID NO:28)
 RRERNRQAAA ANPENSRKG RRQRGKNRG (SEQ ID NO:29)
 P RRERNRQAAA ANPENSRKG RRQRGKNRG (SEQ ID NO:30)
 LP RRERNRQAAA ANPENSRKG RRQRGKNRG (SEQ ID NO:31)
 VLP RRERNRQAAA ANPENSRKG RRQRGKNRG (SEQ ID NO:32)
 AVLPL RRERNRQAAA ANPENSRKG RRQRGKNRG (SEQ ID NO:33)
 MAVLP RRERNRQAAA ANPENSRKG RRQRGKNRG (SEQ ID NO:34)
 QMAVLP RRERNRQAAA ANPENSRKG RRQRGKNRG (SEQ ID NO:35)
 KQMAVLP RRERNRQAAA ANPENSRKG RRQRGKNRG (SEQ ID NO:36)
 DKQMAVLP RRERNRQAAA ANPENSRKG RRQRGKNRG (SEQ ID NO:37)
 PDKQMAVLP RRERNRQAAA ANPENSRKG RRQRGKNRG (SEQ ID NO:38)

As used herein, the term "truncated GDNF protein product" includes biologically active synthetic or recombinant truncated GDNF proteins, truncated GDNF proteins produced from mature GDNF, biologically active truncated GDNF variants (including insertion, substitution and deletion variants), and chemically modified derivatives thereof. Also included are truncated GDNF proteins that are substantially homologous to the human GDNF protein having the amino acid sequence set forth in SEQ ID NO:2.

The term "biologically active" as used herein means that the truncated GDNF protein demonstrates similar neurotrophic properties, but not necessarily all of the same properties, and not necessarily to the same degree, as the GDNF protein having the amino acid sequence set forth in SEQ ID NO:2. The selection of the particular neurotrophic properties of interest depends upon the use for which the truncated GDNF protein product is being administered. The truncated GDNF protein products are biologically active and demonstrate dopaminergic neuron survival characteristics similar to that demonstrated by mature GDNF protein using the evaluation of dopamine uptake and tyrosine hydroxylase (TH) expression as an exemplary bioassay, as discussed in the examples, below.

The term "substantially homologous", as used herein, means a degree of homology to the human GDNF having the amino acid sequence set forth in SEQ ID NO:2 that is preferably in excess of 70%, most preferably in excess of 80%, and even more preferably in excess of 90% or even 95%. The percentage of homology as described herein is calculated as the percentage of amino acid residues found in the smaller of the two sequences which align with identical amino acid residues in the sequence being compared when four gaps in a length of 100 amino acids may be introduced to assist in that alignment as set forth by Dayhoff, in *Atlas of Protein Sequence and Structure* Vol. 5, p. 124 (1972), National Biochemical Research Foundation, Washington, D.C., the disclosure of which is hereby incorporated by reference. Also included as substantially homologous is any truncated GDNF protein which may be isolated by virtue of cross-reactivity with antibodies to the GDNF of SEQ ID NO:2 or whose genes may be isolated through hybridization with the gene or with segments of the gene encoding the GDNF of SEQ ID NO:1.

As will be apparent to those skilled in the art upon reading the present description, substantially homologous proteins will involve one or more deletions from, or additions or substitutions to, the amino acid residues of the truncated GDNF protein represented by X-[Cys⁴¹-Cys¹³³]-Y. The production

of such variants is described in further detail below. It will be further appreciated that because the present invention clearly addresses "truncated" GDNF proteins, the amino-terminus addition variants are contemplated as including the addition of a methionine residue, or non-GDNF amino acid residue or sequence, but do not include the addition of an amino acid residue(s) that would result in the reconstruction of the mature GDNF protein. Truncated GDNF proteins based upon naturally occurring allelic mutants or variants are also within the scope of the present invention. The production of variant truncated GDNF protein is described in further detail below.

Lin et al. (U.S. Patent Application No. 08/182,183, *supra*) described the truncation of mature GDNF at the carboxyl terminus by proteolytic processing of the Lys-Arg residues which are the sixth and fifth residues, respectively, from the carboxyl terminal end of mature GDNF (i.e., Lys¹²⁹-Arg¹³⁰ according to the amino acid residue numbering of Figure 1 (also as in SEQ ID NO:1 or SEQ ID NO:2). Such a truncation would eliminate two cysteine residues from the mature GDNF protein. This would be likely to result in improper folding of the protein, and therefore, would result in the formation of an inactive protein. In contrast, the X-[Cys⁴¹-Cys¹³³]-Y truncated GDNF protein products of the present invention retain the Cys¹³¹ and Cys¹³³ residues and are active proteins as determined by dopamine uptake assay.

In one embodiment of the present invention, preferred truncated GDNF protein products lack one or more deamidation sites. Such a lack of deamidation sites would result in the enhanced biochemical stability of the purified protein and a decrease in possible degradation products, thereby resulting in a more storage stable protein. An exemplary truncated GDNF protein product is the [Ser²⁶-Ile¹³⁴] truncated GDNF protein which lacks the sites which otherwise may lead to deamidation of the mature protein. Alternatively, the [Arg¹⁶-Ile¹³⁴] truncated GDNF protein would lack at least the first deamidation site otherwise present in the mature protein.

A currently preferred truncated GDNF protein product is [Arg³²-Ile¹³⁴] truncated GDNF protein. This truncated GDNF protein lacks the site at or near which proteolytic clipping of the mature protein occurs. Therefore, this truncated GDNF protein is expected to be resistant to the processing event which may also occur *in vivo*. Another currently preferred truncated GDNF protein product is the [Lys³⁷-Ile¹³⁴] truncated GDNF protein. This truncation would further reduce the pI of the truncated protein, as would other truncations in

which residues up to and including Gly⁴⁰ and Ile¹³⁴ are removed from the N- and C-terminals, respectively. The presently most preferred truncated GDNF

protein products retain all of the cysteine residues found in mature GDNF protein, but lack any discernible sites for rapid proteolytic processing of the truncated GDNF protein during expression and manufacturing or following *in vivo* administration. These preferred proteins include the [Arg³²-Ile¹³⁴], [Gly³³-Ile¹³⁴], [Gln³⁴-Ile¹³⁴], [Arg³⁵-Ile¹³⁴], [Gly³⁶-Ile¹³⁴], [Lys³⁷-Ile¹³⁴], [Asn³⁸-Ile¹³⁴] and [Arg³⁹-Ile¹³⁴] truncated GDNF protein products.

Similar to the results previously described for mature GDNF by Lin et al. (U.S. Patent Application No. 07/855,413, *supra*), the truncated GDNF proteins of the present invention have demonstrated the ability to increase dopamine uptake by the embryonic precursors of the substantia nigra dopaminergic neurons. Bioassays of the truncated GDNF proteins are further described in

Example 4, below.

The novel truncated GDNF proteins are typically isolated and purified to form truncated GDNF proteins which are substantially free from the presence of other (non-GDNF) proteinaceous materials. Preferably, the truncated GDNF protein products are about 80% free of other proteins which may be present due to the production technique used in the manufacture of the truncated GDNF protein product. More preferably, the truncated GDNF protein products are about 90% free of other proteins, particularly preferably, about 95% free of other proteins, and most preferably about >98% free of other proteins. In addition, the present invention furnishes the unique advantage of providing polynucleotide sequences for the manufacture of homogeneous truncated GDNF proteins. For example, the use of the polynucleotide sequence encoding the [Arg³²-Ile¹³⁴] truncated GDNF protein allows the recombinant production of the truncated GDNF protein in *E. coli* and other appropriate expression systems. In other words, the novel polynucleotides allow the production of truncated GDNF proteins which are not susceptible to proteolytic processing, or which have reduced susceptibility to such processing or other biochemical processing effects as described above. Thus, the novel polynucleotides make it easier to prepare and/or isolate single species truncated GDNF proteins, and therefore, the truncated GDNF proteins and/or products thereof do not contain or contain decreased amounts of the above-described mixture of hetero- and homodimers. It will be appreciated, however, that the final truncated GDNF protein products

may be combined with other factors, chemical compositions and/or suitable pharmaceutical formulation materials prior to administration, as described in further detail below.

- 5 In one aspect of the present invention, the truncated GDNF proteins are advantageously produced via recombinant techniques because they are capable of achieving comparatively higher amounts of protein at greater purity. Recombinant truncated GDNF protein forms include glycosylated and non-glycosylated forms of the protein, and protein expressed in bacterial, mammalian or insect cell systems. Alternatively, the truncated GDNF proteins
10 may be chemically synthesized. Currently preferred production methods are described in greater detail below.

Truncated GDNF Variants and Derivatives

A. Truncated GDNF Variants

- 15 Another aspect of the present invention includes variants of truncated GDNF protein. The term "truncated GDNF protein products" as used herein includes variant proteins in which amino acids have been deleted from ("deletion variants"), inserted into ("addition variants"), or substituted for ("substitution variants"), residues within the amino acid sequence of naturally-occurring
20 GDNF. Such variants are prepared by introducing appropriate nucleotide changes into the DNA encoding the protein or by *in vitro* chemical synthesis of the desired protein. It will be appreciated by those skilled in the art that many combinations of deletions, insertions, and substitutions can be made provided that the final protein possesses GDNF biological activity.
- 25 Mutagenesis techniques for the replacement, insertion or deletion of one or more selected amino acid residues are well known to one skilled in the art (e.g., U.S. Pat. No. 4,518,584, the disclosure of which is hereby incorporated by reference.) There are two principal variables in the construction of amino acid sequence variants: the location of the mutation site and the nature of the
30 mutation. In designing truncated GDNF variants, the location of the mutation site and the nature of the mutation will depend on the biochemical characteristic(s) to be modified. The mutation sites can be modified individually or in series, e.g., by (1) substituting first with conservative amino acid choices and then with more radical selections depending upon the results achieved, (2) deleting the
35 target amino acid residue, or (3) inserting amino acid residues adjacent to the located site.

Amino acid sequence deletions generally range from about 1 to 30 amino acid residues, more usually from about 1 to 10 residues, and typically from about 1 to 5 residues. For example, deletions in the "X" portion of the amino acid residues located N-terminally to Cys⁴¹ may range from approximately 1 to 30 residues, while deletions between the cysteine residues of [Cys⁴¹-Cys¹³³] are typically from about 1 to 5 residues, depending on the location, so as not to disrupt protein folding. Deletions within the truncated GDNF proteins may be made in regions of low homology with transforming growth factor-beta (TGF- β) family members. Deletions from truncated GDNF proteins in areas of substantial homology with other TGF- β family sequences will be more likely to modify the biological activity more significantly. The number of total deletions and/or consecutive deletions will be selected so as to preserve the tertiary structure of truncated GDNF protein in the affected domain, e.g., cysteine crosslinking.

Amino acid sequence additions may include amino- and/or carboxyl-terminal fusions ranging in length from one residue to one hundred or more residues, as well as internal intrasequence insertions of single or multiple amino acid residues. Internal additions may range generally from about 1 to 10 amino acid residues, more typically from about 1 to 5 amino acid residues, and usually from about 1 to 3 amino acid residues. As described above, the amino-terminus addition variants of the present invention are contemplated as including the addition of a methionine (for example, as an artifact of the direct expression of GDNF in bacterial recombinant cell culture) or a non-GDNF amino acid residue or sequence. Amino-terminus addition variants do not involve the addition of an amino acid residue(s) that would result in the reconstruction of the mature GDNF protein. A further example of a terminal insertion includes the fusion of a heterologous N-terminal signal sequence to the N-terminus to facilitate the secretion of protein from recombinant host cells. Such signal sequences generally will be obtained from, and thus be homologous to, the intended host cell species. Insertions or additions may also include amino acid sequences derived from the sequence of other neurotrophic factors.

Another group of variants are amino acid substitution variants. These variants have at least one amino acid residue in the truncated GDNF protein removed and a different residue inserted in its place. See, for example, Figure 5 wherein naturally occurring Asn²² was changed to Ser to facilitate further removal of the Met residue. Using the X-[Cys⁴¹-Cys¹³³]-Y amino acid sequence representation and the present definition of truncated GDNF protein

products, such a truncated GDNF protein may be referred to either as a substitution variant Met-[Asn²²ΔSer²²-Ile¹³⁴] truncated GDNF protein or an addition variant Met-Ser-[Pro²³-Ile¹³⁴] truncated GDNF protein.

Substitution variants include allelic variants, which are characterized by naturally-occurring nucleotide sequence changes in the species population that may or may not result in an amino acid change.

Specific mutations of the sequences of the truncated GDNF proteins may involve modifications of a glycosylation site (e.g., serine, threonine, or asparagine). The absence of glycosylation or only partial glycosylation may result from amino acid substitution or deletion at any asparagine-linked glycosylation recognition site or at any site of the protein that is modified by the addition of an O-linked carbohydrate. An asparagine-linked glycosylation recognition site comprises a tripeptide sequence which is specifically recognized by appropriate cellular glycosylation enzymes. These tripeptide sequences are either Asn-Xaa-Thr or Asn-Xaa-Ser, where Xaa can be any amino acid other than Pro. A variety of amino acid substitutions or deletions at one or both of the first or third amino acid positions of a glycosylation recognition site (and/or amino acid deletion at the second position) result in non-glycosylation at the modified tripeptide sequence. Thus, the expression of appropriately altered nucleotide sequences produces variants which are not glycosylated at that site. Alternatively, the sequence may be modified to add glycosylation sites to the truncated GDNF protein.

One method for identifying truncated GDNF amino acid residues or regions for mutagenesis is called "alanine scanning mutagenesis" as described by Cunningham and Wells (*Science*, 244: 1081-1085, 1989). In this method, an amino acid residue or group of target residues are identified (e.g., charged residues such as Arg, Asp, His, Lys, and Glu) and replaced by a neutral or negatively charged amino acid (most preferably alanine or polyalanine) to effect the interaction of the amino acids with the surrounding aqueous environment in or outside the cell. Those domains demonstrating functional sensitivity to the substitutions then are refined by introducing additional or alternate residues at the sites of substitution. Thus, the site for introducing an amino acid sequence modification is predetermined, and to optimize the performance of a mutation at a given site, alanine scanning or random mutagenesis may be conducted and the variants are screened for the optimal combination of desired activity and degree of activity.

The sites of greatest interest for substitutional mutagenesis include sites where the amino acids found in GDNF proteins from various species are substantially different in terms of side-chain bulk, charge, and/or hydrophobicity. Other sites of interest include those in which particular residues of GDNF-like proteins, obtained from various species, are identical. Such positions are generally important for the biological activity of a protein. Initially, these sites are modified by substitution in a relatively conservative manner. Such conservative substitutions are shown in Table 1 under the heading of preferred substitutions. If such substitutions result in a change in biological activity, then more substantial changes (exemplary substitutions) are introduced and/or other additions/deletions may be made, and the resulting products screened.

TABLE 1
Amino Acid Substitutions

<u>Original Residue</u>	<u>Preferred Substitutions</u>	<u>Exemplary Substitutions</u>
Ala (A)	Val	Val; Leu; Ile
Arg (R)	Lys	Lys; Gln; Asn
Asn (N)	Gln	Gln; His; Lys; Arg
Asp (D)	Glu	Glu
Cys (C)	Ser	Ser
Gln (Q)	Asn	Asn
Glu (E)	Asp	Asp
Gly (G)	Pro	Pro
His (H)	Arg	Asn; Gln; Lys; Arg
Ile (I)	Leu	Leu; Val; Met; Ala; Phe; norleucine
Leu (L)	Ile	norleucine; Ile; Val; Met; Ala; Phe
Lys (K)	Arg	Arg; Gln; Asn
Met (M)	Leu	Leu; Phe; Ile
Phe (F)	Leu	Leu; Val; Ile; Ala
Pro (P)	Gly	Gly
Ser (S)	Thr	Thr
Thr (T)	Ser	Ser
Trp (W)	Tyr	Tyr
Tyr (Y)	Phe	Trp; Phe; Thr; Ser
Val (V)	Leu	Ile; Leu; Met; Phe; Ala; norleucine

Conservative modifications to the amino acid sequence (and the corresponding modifications to the encoding nucleic acid sequences) are expected

to produce truncated GDNF proteins having functional and chemical characteristics similar to those of the truncated GDNF proteins described in the Examples, below. In contrast, substantial modifications in the functional and/or chemical characteristics of truncated GDNF proteins may be accomplished by

- 5 selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the protein at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side chain
- 10 properties:

- 1) hydrophobic: norleucine, Met, Ala, Val, Leu, Ile;
- 2) neutral hydrophilic: Cys, Ser, Thr;
- 3) acidic: Asp, Glu;
- 4) basic: Asn, Gln, His, Lys, Arg;
- 15 5) residues that influence chain orientation: Gly, Pro; and
- 6) aromatic: Trp, Tyr, Phe.

Non-conservative substitutions may involve the exchange of a member of one of these classes for another. Such substituted residues may be introduced into regions of the truncated GDNF proteins that are homologous with other TGF-
20 β proteins, or into the non-homologous regions of the protein.

B. Truncated GDNF Derivatives

Chemically modified derivatives of truncated GDNF or truncated GDNF variants may be prepared by one skilled in the art given the disclosures herein.

- 25 The chemical moieties most suitable for derivatization of truncated GDNF proteins include water soluble polymers. A water soluble polymer is desirable because the protein to which it is attached does not precipitate in an aqueous environment, such as a physiological environment. Preferably, the polymer will be pharmaceutically acceptable for the preparation of a therapeutic product or composition. One skilled in the art will be able to select the desired polymer
- 30 based on such considerations as whether the polymer/protein conjugate will be used therapeutically, and if so, the desired dosage, circulation time, resistance to proteolysis, and other considerations. The effectiveness of the derivatization may be ascertained by administering the derivative, in the desired form (i.e., by
- 35 osmotic pump, or, more preferably, by injection or infusion, or further

formulated for oral, pulmonary or other delivery routes), and determining its effectiveness.

Suitable water soluble polymers include, but are not limited to, polyethylene glycol (PEG), copolymers of ethylene glycol/propylene glycol, monomethoxy-polyethylene glycol, carboxymethylcellulose, dextran, polyvinyl alcohol, polyvinyl pyrrolidone, poly-1, 3-dioxolane, poly-1,3,6-trioxane, ethylene/maleic anhydride copolymer, polyaminoacids (either homopolymers or random copolymers), poly(n-vinyl pyrrolidone)polyethylene glycol, propylene glycol homopolymers, polypropylene oxide/ethylene oxide co-polymers, polyoxyethylated polyols (e.g., glycerol), polyethylene glycol propionaldehyde, and mixtures thereof. As used herein, polyethylene glycol is meant to encompass any of the forms of PEG that have been used to derivatize other proteins, such as mono-(C1-C10) alkoxy- or aryloxy-polyethylene glycol. Polyethylene glycol propionaldehyde may have advantages in manufacturing due to its stability in water. The polymer may be of any molecular weight, and may be branched or unbranched.

The present invention particularly relates to truncated GDNF protein products involving truncated GDNF protein linked to at least one PEG molecule. In another aspect, the present invention relates to truncated GDNF protein attached to at least one PEG molecule via an acyl or alkyl linkage.

Pegylation may be carried out by any of the pegylation reactions known in the art. See, for example: *Focus on Growth Factors* 3(2): 4-10 (1992); EP 0 154 316; EP 0 401 384; and Malik et al., *Exp. Hematol.* 20: 1028-1035 (1992) (reporting pegylation of GM-CSF using tresyl chloride). Preferably, the pegylation is carried out via an acylation reaction or an alkylation reaction with a reactive water soluble polymer. These preferred means for derivatization are discussed in greater detail, below. For the acylation reactions, the polymer(s) selected preferably have a single reactive ester group. For the reductive alkylation reactions, the polymer(s) selected preferably have a single reactive aldehyde group. In addition, the selected polymer may be modified to have a single reactive group, such as an active ester for acylation or an aldehyde for alkylation, so that the degree of polymerization may be controlled. Generally, the water soluble polymer will not be selected from naturally-occurring glycosyl residues since these are usually made more conveniently by mammalian recombinant expression systems.

Acylation

In the present invention, pegylation by acylation generally involves reacting an active ester derivative of polyethylene glycol with a truncated GDNF protein. Any known or subsequently discovered reactive PEG molecule may be used to carry out the pegylation process. A preferred activated PEG ester is PEG esterified to N-hydroxysuccinimide ("NHS"). As used herein, "acylation" is contemplated to include without limitation the following types of linkages between a truncated GDNF protein and a water soluble polymer such as PEG: amide, carbamate, urethane, and the like. See *Bioconjugate Chem.* 5: 133-140 (1994). Reaction conditions may be selected from any of those known in the pegylation art or those subsequently developed, but should avoid or limit exposure to reaction conditions such as temperatures, solvents, and pH levels that would inactivate the truncated GDNF protein to be modified.

Pegylation by acylation will generally result in a poly-pegylated truncated GDNF protein, wherein the lysine ϵ -amino groups are pegylated via an acyl linking group. Preferably, the connecting linkage will be an amide. Also preferably, the resulting product will be substantially only (e.g., $\geq 95\%$) mono-, di- or tri- pegylated. However, some conjugates with higher degrees of pegylation may be formed in amounts depending on the specific reaction conditions used. If desired, more purified pegylated conjugates may be prepared from the mixture by standard purification techniques, including, among others, dialysis, salting-out, ultrafiltration, ion-exchange chromatography, gel filtration chromatography and electrophoresis.

Alkylation

In the present invention, pegylation by alkylation generally involves reacting a terminal aldehyde derivative of PEG with a truncated GDNF protein in the presence of a reducing agent. Pegylation by alkylation can also result in a poly-pegylated truncated GDNF protein. In addition, one can manipulate the reaction conditions to favor pegylation substantially only at the α -amino group of the N-terminus of the protein (i.e., a mono-pegylated species). In either case of mono-pegylation or polypegylation, the PEG groups are preferably attached to the protein via a $-\text{CH}_2\text{-NH}-$ group. With particular reference to the $-\text{CH}_2-$ group, this type of linkage is referred to herein as an "alkyl" linkage.

Selective N-terminal chemical modification may be accomplished by reductive alkylation which exploits differential reactivity of different types of

primary amino groups (lysine versus the N-terminal) available for derivatization in a particular protein. Under the appropriate reaction conditions, substantially selective derivatization of the protein at the N-terminus with a carbonyl group-containing polymer is achieved. For example, one may selectively N-terminally pegylate the protein by performing the reaction at a pH which allows one to take advantage of the pK_a differences between the ϵ -amino group of the lysine residues and that of the α -amino group of the N-terminal residue of the protein. By such selective derivatization, attachment of a water soluble polymer to a protein is controlled: the conjugation with the polymer takes place predominantly at the N-terminus of the protein and no significant modification of other reactive groups, such as the lysine side chain amino groups, occurs. Using reductive alkylation, the water soluble polymer preferably has a single reactive aldehyde for coupling to the protein. Polyethylene glycol propionaldehyde, containing a single reactive aldehyde, may be used.

The present invention includes pegylated truncated GDNF proteins, wherein the PEG group(s) is (are) attached via acyl or alkyl groups. As discussed above, such truncated GDNF protein products may be mono-pegylated or poly-pegylated (e.g., containing 2-6, preferably 2-5, PEG groups). The PEG groups are generally attached to the protein at the α - or ϵ -amino groups of amino acids, but it is also contemplated that the PEG groups could be attached to any reactive group of the protein, which is sufficiently reactive to become attached to a PEG group under suitable reaction conditions. Thus, polyethylene glycol may be covalently bound to a protein via a reactive group, such as, a free amino or carboxyl group. Reactive groups are those to which an activated PEG molecule may be bound. The amino acid residues having a free amino group may include lysine residues and the N-terminal amino acid residue. Those having a free carboxyl group may include aspartic acid residues, glutamic acid residues, and the C-terminal amino acid residue. Sulfhydryl groups may also be used as a reactive group for attaching PEG molecule(s). For therapeutic purposes, attachment at an amino group, such as attachment at the N-terminus or lysine group is typically preferred. Attachment at residues important for receptor binding should be avoided if receptor binding is desired.

In one aspect, the present invention provides for a substantially homogeneous preparation of mono-polymer/truncated GDNF protein conjugate

wherein a polymer molecule has been attached substantially only (i.e., $\geq 95\%$) in a single location. More specifically, if PEG is used, the present invention also provides for pegylated a truncated GDNF protein lacking possibly antigenic linking groups, and having the PEG molecule directly coupled to the truncated

5 GDNF protein.

In addition, derivatives may be prepared using glycosylated, non-glycosylated or de-glycosylated truncated GDNF proteins. Typically, non-glycosylated truncated GDNF proteins are used. For example, the prokaryote-expressed [Arg³²-Ile¹³⁴] truncated GDNF protein may be chemically

10 derivatized to include mono- or poly-, e.g., 2-4, PEG moieties, attached via an acyl or alkyl group):

In general, chemical derivatization may be performed under any suitable condition used to react a biologically active substance with an activated polymer molecule. Methods for preparing pegylated truncated GDNF proteins will

15 generally comprise the steps of (a) reacting a truncated GDNF protein with polyethylene glycol (such as a reactive ester or aldehyde derivative of PEG) under conditions whereby the truncated GDNF protein becomes attached to one or more PEG groups, and (b) obtaining the reaction product(s). In general, the optimal reaction conditions for the acylation reactions will be determined case-

20 by-case based on known parameters and the desired result. For example, the larger the ratio of PEG:protein, the greater the percentage of poly-pegylated product. The optimum ratio (in terms of efficiency of reaction in that there is no excess unreacted protein or polymer) may be determined by factors such as the desired degree of derivatization (e.g., mono-, di-, tri-, etc.), the molecular

25 weight of the polymer selected, whether the polymer is branched or unbranched, and the reaction conditions used.

Reductive alkylation to produce a substantially homogeneous population of mono-polymer/truncated GDNF protein conjugate will generally comprise the steps of (a) reacting a truncated GDNF protein with a reactive PEG molecule

30 under reductive alkylation conditions, at a pH suitable to permit selective modification of the α -amino group at the amino terminus of the truncated GDNF protein, and (b) obtaining the reaction product(s).

For a substantially homogeneous population of mono-polymer/truncated GDNF protein conjugate, the reductive alkylation reaction conditions are those

35 which permit the selective attachment of the water soluble polymer moiety to the N-terminus of a truncated GDNF protein. Such reaction conditions generally

provide for pK_a differences between the lysine amino groups and the α -amino group at the N-terminus (the pK_a being the pH at which 50% of the amino groups are protonated and 50% are not). The pH also affects the ratio of polymer to protein to be used. In general, if the pH is lower, a larger excess of polymer to protein will be desired (i.e., the less reactive the N-terminal α -amino group, the more polymer needed to achieve optimal conditions). If the pH is higher, the polymer:protein ratio need not be as large (i.e., more reactive groups are available, so fewer polymer molecules are needed). For purposes of the present invention, the pH will generally fall within the range of 3-9, preferably 3-6.

Another consideration is the molecular weight of the polymer. In general, the higher the molecular weight of the polymer, the fewer the number of polymer molecules which may be attached to the protein. Similarly, branching of the polymer should be taken into account when optimizing these parameters. Generally, the higher the molecular weight (or the more branches) the higher the polymer:protein ratio. In general, for the pegylation reactions contemplated herein, the preferred average molecular weight is about 2 kDa to about 100 kDa (the term "about" indicating that in preparations of polyethylene glycol, some molecules will weigh more, some less, than the stated molecular weight). The preferred average molecular weight is about 5 kDa to about 50 kDa, particularly preferably about 12 kDa to about 25 kDa. The ratio of water-soluble polymer to truncated GDNF protein will generally range from 1:1 to 100:1, preferably (for polypegylation) 1:1 to 20:1, and (for mono-pegylation) 1:1 to 5:1.

Using the conditions indicated above, reductive alkylation will provide for selective attachment of the polymer to any truncated GDNF protein having an α -amino group at the amino terminus, and provide for a substantially homogenous preparation of mono-polymer/truncated GDNF protein conjugate. The term "mono-polymer/truncated GDNF protein conjugate" is used here to mean a derivative containing a single polymer molecule attached to a truncated GDNF protein. The mono-polymer/truncated GDNF protein conjugate preferably will have a polymer molecule located at the N-terminus, but not on lysine amino side groups. The preparation will preferably be greater than 90% mono-polymer/truncated GDNF protein conjugate, and more preferably greater than 95% mono-polymer/truncated GDNF protein conjugate, with the remainder of observable proteins being unreacted (i.e., protein lacking the polymer moiety).

For reductive alkylation, the reducing agent should be stable in aqueous solution and preferably be able to reduce only the Schiff base formed in the

initial process of reductive alkylation. Exemplary reducing agents may be selected from the group consisting of sodium borohydride, sodium cyanoborohydride, dimethylamine borane, trimethylamine borane and pyridine borane. A particularly preferred reducing agent is sodium cyanoborohydride.

- 5 Other reaction parameters, such as solvent, reaction times, temperatures, etc., and means of purification of products, can be determined case-by-case based on commonly available information relating to derivatization of proteins with water soluble polymers.

- 10 One may choose to prepare a mixture of polymer/protein conjugates by acylation and/or alkylation methods, and the advantage provided herein is that one may select the proportion of mono-polymer/ protein conjugate to include in the mixture. Thus, if desired, one may prepare a mixture of protein having various numbers of polymer molecules attached thereto (i.e., di-, tri-, tetra-, etc.) and combine with the mono-polymer/protein conjugate material prepared
- 15 using the present methods, and have a mixture with a predetermined proportion of mono-polymer/protein conjugate.

Polynucleotides Encoding Truncated GDNF Proteins

- The present invention further provides novel polynucleotides which
- 20 encode truncated GDNF proteins. When used as a hybridization probe or amplification primer, the nucleic acid sequence will be substantially free from all other nucleic acid sequences. For use in recombinant protein expression, the nucleic acid sequence will generally be substantially free from nucleic acid sequences encoding other proteins, unless a fusion protein is intended. Based
- 25 upon the present description and using the universal codon table, one of ordinary skill in the art can readily determine all of the nucleic acid sequences which encode the amino acid sequences of truncated GDNF proteins. Presently preferred nucleic acid sequences include those polynucleotides encoding the [Arg¹⁶-Ile¹³⁴], [Ser²⁶-Ile¹³⁴], [Arg³²-Ile¹³⁴], and [Lys³⁷-Ile¹³⁴] truncated
- 30 GDNF proteins. Examples of a variety of polynucleotides are depicted in Figures 5, 6 and 7 as well as those portions of Figures 1, 3 and 4 which encode truncated GDNF proteins. It will also be appreciated by those skilled in the art that the novel polynucleotides which encode truncated GDNF proteins include those nucleic acid sequences encoding variant truncated GDNF proteins, whether man-
- 35 made or naturally occurring.

Recombinant expression techniques, conducted in accordance with the descriptions set forth below, may be followed to produce these polynucleotides and express the various truncated GDNF proteins. For example, by inserting a nucleic acid sequence which encodes a truncated GDNF protein into an appropriate vector, one skilled in the art can readily produce large quantities of the desired nucleotide sequence. The sequences can then be used to generate detection probes or amplification primers. Alternatively, a polynucleotide encoding a truncated GDNF protein can be inserted into an expression vector. By introducing the expression vector into an appropriate host, the desired truncated GDNF protein may be produced in large amounts.

As further described herein, there are numerous host/vector systems available for the propagation of nucleic acid sequences and/or the production of truncated GDNF proteins. These include, but are not limited to, plasmid, viral and insertional vectors, and prokaryotic and eukaryotic hosts. One skilled in the art can adapt a host/vector system which is capable of propagating or expressing heterologous DNA to produce or express the sequences of the present invention.

By means of such recombinant techniques, the truncated GDNF proteins of the present invention are readily produced in commercial quantities. Furthermore, it will be appreciated by those skilled in the art that, in view of the present disclosure, the novel nucleic acid sequences include degenerate nucleic acid sequences encoding the truncated GDNF proteins specifically set forth in the Figures, variants of such truncated GDNF proteins, and those nucleic acid sequences which hybridize, preferably under stringent hybridization conditions, to complements of these nucleic acid sequences (see, Maniatis et. al., *Molecular Cloning* (A Laboratory Manual); Cold Spring Harbor Laboratory, pages 387 to 389, 1982.) Exemplary stringent hybridization conditions are hybridization in 4 x SSC at 62-67°C, followed by washing in 0.1 x SSC at 62-67°C for approximately an hour. Alternatively, exemplary stringent hybridization conditions are hybridization in 45-55% formamide, 4 x SSC at 40-45°C. DNA sequences which hybridize to the complementary sequences for truncated GDNF protein under relaxed hybridization conditions and which encode a truncated GDNF protein of the present invention are also included herein. Examples of such relaxed stringency hybridization conditions are 4 x SSC at 45-55°C or hybridization with 30-40% formamide at 40-45°C.

Also provided by the present invention are recombinant DNA constructs involving vector DNA together with the DNA sequence encoding a truncated GDNF

protein. In such DNA constructs, the nucleic acid sequence encoding truncated GDNF protein (with or without signal peptides) is in operative association with a suitable expression control or regulatory sequence capable of directing the replication and/or expression of the truncated GDNF protein in a selected host.

5

Recombinant Expression of Truncated GDNF Protein

Preparation of Polynucleotides encoding Truncated GDNF

A nucleic acid sequence encoding truncated GDNF, or a mature GDNF starting material, can readily be obtained in a variety of ways, including, without limitation, chemical synthesis, cDNA or genomic library screening, expression library screening, and/or PCR amplification of cDNA. These methods and others useful for isolating such nucleic acid sequences are set forth, for example, by Sambrook *et al.* (*Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989), by Ausubel *et al.*, eds (*Current Protocols in Molecular Biology*, Current Protocols Press, 1994), and by Berger and Kimmel (*Methods in Enzymology: Guide to Molecular Cloning Techniques*, vol. 152, Academic Press, Inc., San Diego, CA, 1987). Preferred nucleic acid sequences encoding GDNF are mammalian sequences.

Chemical synthesis of a nucleic acid sequence which encodes a truncated GDNF protein can also be accomplished using methods well known in the art, such as those set forth by Engels *et al.* (*Angew. Chem. Intl. Ed.*, 28:716-734, 1989). These methods include, *inter alia*, the phosphotriester, phosphoramidite and H-phosphonate methods of nucleic acid sequence synthesis. The nucleic acid sequence encoding the truncated GDNF protein will be several hundred base pairs (bp) or nucleotides in length. Large nucleic acid sequences, for example those larger than about 100 nucleotides in length, can be synthesized as several fragments. The fragments can then be ligated together to form a nucleic acid sequence encoding truncated GDNF protein. A preferred method is polymer-supported synthesis using standard phosphoramidite chemistry.

Alternatively, a suitable nucleic acid sequence may be obtained by screening an appropriate cDNA library (i.e., a library prepared from one or more tissue source(s) believed to express the protein) or a genomic library (a library prepared from total genomic DNA). The source of the cDNA library is typically a tissue from any species that is believed to express GDNF in reasonable quantities. The source of the genomic library may be any tissue or tissues from any mammalian or other species believed to harbor a gene encoding

GDNF or a GDNF homologue. The library can be screened for the presence of the GDNF cDNA/gene using one or more nucleic acid probes (oligonucleotides, cDNA or genomic DNA fragments that possess an acceptable level of homology to the GDNF or GDNF homologue cDNA or gene to be cloned) that will hybridize selectively with GDNF or GDNF homologue cDNA(s) or gene(s) present in the library. The probes typically used for such library screening usually encode a small region of the GDNF DNA sequence from the same or a similar species as the species from which the library was prepared. Alternatively, the probes may be degenerate, as discussed herein.

Library screening is typically accomplished by annealing the oligonucleotide probe or cDNA to the clones in the library under conditions of stringency that prevent non-specific binding but permit binding of those clones that have a significant level of homology with the probe or primer. Typical hybridization and washing stringency conditions depend in part on the size (i.e., number of nucleotides in length) of the cDNA or oligonucleotide probe, and whether the probe is degenerate. The probability of obtaining a clone(s) is also considered in designing the hybridization solution (i.e., whether a cDNA or genomic library is being screened; if it is a cDNA library, the probability that the cDNA of interest is present at a high level).

Where DNA fragments (such as cDNAs) are used as probes, typical hybridization conditions include those as set forth in Ausubel et al., *eds., supra*. After hybridization, the blot containing the library is washed at a suitable stringency, depending on several factors such as probe size, expected homology of probe to clone, type of library being screened, number of clones being screened, and the like. Examples of stringent washing solutions (which are usually low in ionic strength and are used at relatively high temperatures) are as follows. One such stringent wash is 0.015 M NaCl, 0.005 M NaCitrate and 0.1% SDS at 55-65°C. Another such stringent buffer is 1 mM Na₂EDTA, 40 mM NaHPO₄, pH 7.2, and 1% SDS at about 40-50°C. One other stringent wash is 0.2 X SSC and 0.1% SDS at about 50-65°C.

There are also exemplary protocols for stringent washing conditions where oligonucleotide probes are used to screen cDNA or genomic libraries. For example, a first protocol uses 6 X SSC with 0.05 percent sodium pyrophosphate at a temperature of between about 35 and 62°C, depending on the length of the probe. For example, 14 base probes are washed at 35-40°C, 17 base probes at 45-50°C, 20 base probes at 52-57°C, and 23 base probes at 57-63°C. The

temperature can be increased 2-3°C where the background non-specific binding appears high. A second protocol uses tetramethylammonium chloride (TMAC) for washing. One such stringent washing solution is 3 M TMAC, 50 mM Tris-HCl, pH 8.0, and 0.2% SDS.

5 Another suitable method for obtaining a nucleic acid sequence encoding a GDNF protein is the polymerase chain reaction (PCR). In this method, poly(A)+RNA or total RNA is extracted from a tissue that expresses GDNF. cDNA is then prepared from the RNA using the enzyme reverse transcriptase. Two primers, typically complementary to two separate regions of the GDNF cDNA
10 (oligonucleotides), are then added to the cDNA along with a polymerase such as *Taq* polymerase, and the polymerase amplifies the cDNA region between the two primers.

Where the method of choice for preparing the nucleic acid sequence encoding the desired truncated GDNF protein requires the use of oligonucleotide
15 primers or probes (e.g., PCR, cDNA or genomic library screening), the oligonucleotide sequences selected as probes or primers should be of adequate length and sufficiently unambiguous so as to minimize the amount of non-specific binding that will occur during library screening or PCR amplification. The actual sequence of the probes or primers is usually based on conserved or highly
20 homologous sequences or regions from the same or a similar gene from another organism. Optionally, the probes or primers can be fully or partially degenerate, i.e., contain a mixture of probes/primers, all encoding the same amino acid sequence, but using different codons to do so. An alternative to
25 preparing degenerate probes is to place an inosine in some or all of those codon positions that vary by species. The oligonucleotide probes or primers may be prepared by chemical synthesis methods for DNA as described above.

Truncated GDNF proteins based on these nucleic acid sequences encoding GDNF, as well as mutant or variant sequences thereof, are also contemplated as within the scope of the present invention. As described above, a mutant or
30 variant sequence is a sequence that contains one or more nucleotide substitutions, deletions, and/or insertions as compared to the wild type sequence and that results in the expression of amino acid sequence variations as compared to the wild type amino acid sequence. In some cases, naturally occurring GDNF amino acid mutants or variants may exist, due to the existence of natural allelic
35 variation. Truncated GDNF proteins based on such naturally occurring mutants or variants are also within the scope of the present invention. Preparation of

synthetic mutant sequences is also well known in the art, and is described for example in Wells et al. (*Gene*, 34:315, 1985) and in Sambrook et al., *supra*.

Vectors

5 The cDNA or genomic DNA encoding a truncated GDNF protein is inserted into a vector for further cloning (amplification of the DNA) or for expression. Suitable vectors are commercially available, or the vector may be specially constructed. The selection or construction of the appropriate vector will depend on 1) whether it is to be used for DNA amplification or for DNA expression, 2) 10 the size of the DNA to be inserted into the vector, and 3) the host cell (e.g., mammalian, insect, yeast, fungal, plant or bacterial cells) to be transformed with the vector. Each vector contains various components depending on its function (amplification of DNA or expression of DNA) and its compatibility with the intended host cell. The vector components generally include, but are not 15 limited to, one or more of the following: a signal sequence, an origin of replication, one or more selection or marker genes, enhancer elements, promoters, a transcription termination sequence, and the like. These components may be obtained from natural sources or synthesized by known procedures. The vectors of the present invention involve a nucleic acid sequence 20 which encodes the truncated GDNF protein of interest operatively linked to one or more of the following expression control or regulatory sequences capable of directing, controlling or otherwise effecting the expression of the truncated GDNF protein by a selected host cell.

25 Signal Sequence

The signal sequence may be a component of the vector, or it may be a part of GDNF DNA that is inserted into the vector. The native GDNF DNA encodes a signal sequence at the amino terminus of the protein that is cleaved during post-translational processing of the protein to form the mature GDNF protein. 30 Included within the scope of this invention are truncated GDNF polynucleotides with the native signal sequence and other pre-pro sequences as well as truncated GDNF polynucleotides wherein the native signal sequence is deleted and replaced with a heterologous signal sequence. The heterologous signal sequence selected should be one that is recognized and processed, i.e., cleaved by a signal peptidase, 35 by the host cell. For prokaryotic host cells that do not recognize and process the native GDNF signal sequence, the signal sequence is substituted by a prokaryotic

signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, or heat-stable enterotoxin II leaders. For yeast secretion, the native GDNF signal sequence may be substituted by the yeast invertase, alpha factor, or acid phosphatase leaders. In mammalian cell expression the native signal sequence is satisfactory, although other mammalian signal sequences may be suitable.

Origin of Replication

Expression and cloning vectors generally include a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. In cloning vectors, this sequence is typically one that enables the vector to replicate independently of the host chromosomal DNA, and includes origins of replication or autonomously replicating sequences. Such sequences are well known for a variety of bacteria, yeasts, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria and various origins (e.g., SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian expression vectors (for example, the SV40 origin is often used only because it contains the early promoter).

Selection Gene

The expression and cloning vectors typically contain a selection gene. This gene encodes a "marker" protein necessary for the survival or growth of the transformed host cells when grown in a selective culture medium. Host cells that were not transformed with the vector will not contain the selection gene, and therefore, they will not survive in the culture medium. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline; (b) complement auxotrophic deficiencies; or (c) supply critical nutrients not available from the culture medium.

Other selection genes may be used to amplify the gene which will be expressed. Amplification is the process wherein genes which are in greater demand for the production of a protein critical for growth are reiterated in tandem within the chromosomes of successive generations of recombinant cells.

Examples of suitable selectable markers for mammalian cells include dihydrofolate reductase (DHFR) and thymidine kinase. The mammalian cell

transformants are placed under selection pressure which only the transformants are uniquely adapted to survive by virtue of the marker present in the vector. Selection pressure is imposed by culturing the transformed cells under conditions in which the concentration of selection agent in the medium is successively changed, thereby leading to amplification of both the selection gene and the DNA that encodes truncated GDNF. As a result, increased quantities of truncated GDNF are synthesized from the amplified DNA.

For example, cells transformed with the DHFR selection gene are first identified by culturing all of the transformants in a culture medium that contains methotrexate, a competitive antagonist of DHFR. An appropriate host cell when wild-type DHFR is used is the Chinese hamster ovary cell line deficient in DHFR activity (see, for example, Urlaub and Chasin, *Proc. Natl. Acad. Sci., USA* 77(7): 4216-4220 (1980)). The transformed cells are then exposed to increased levels of methotrexate. This leads to the synthesis of multiple copies of the DHFR gene, and, concomitantly, multiple copies of other DNA present in the expression vector, such as the DNA encoding a truncated GDNF protein.

Promoter

The expression and cloning vectors of the present invention will typically contain a promoter that is recognized by the host organism and operably linked to the nucleic acid sequence encoding the truncated GDNF protein. Promoters are untranslated sequences located upstream (5') to the start codon of a structural gene (generally within about 100 to 1000 bp) that control the transcription and translation of a particular nucleic acid sequence, such as that encoding truncated GDNF. Promoters are conventionally grouped into one of two classes, inducible promoters and constitutive promoters. Inducible promoters initiate increased levels of transcription from DNA under their control in response to some change in culture conditions, such as the presence or absence of a nutrient or a change in temperature. A large number of promoters, recognized by a variety of potential host cells, are well known. These promoters are operably linked to the DNA encoding truncated GDNF by removing the promoter from the source DNA by restriction enzyme digestion and inserting the desired promoter sequence into the vector. The native GDNF promoter sequence may be used to direct amplification and/or expression of truncated GDNF DNA. A heterologous promoter is preferred, however, if it permits greater transcription and higher

yields of the expressed protein as compared to the native promoter, and if it is compatible with the host cell system that has been selected for use.

Promoters suitable for use with prokaryotic hosts include the beta-lactamase and lactose promoter systems; alkaline phosphatase, a tryptophan (trp) promoter system; and hybrid promoters such as the tac promoter. Other known bacterial promoters are also suitable. Their nucleotide sequences have been published, thereby enabling one skilled in the art to ligate them to the desired DNA sequence(s), using linkers or adaptors as needed to supply any required restriction sites.

Suitable promoting sequences for use with yeast hosts are also well known in the art. Yeast enhancers are advantageously used with yeast promoters. Suitable promoters for use with mammalian host cells are well known and include those obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and most preferably Simian Virus 40 (SV40). Other suitable mammalian promoters include heterologous mammalian promoters, e.g., heat-shock promoters and the actin promoter. A currently used promoter in the production of GDNF proteins in CHO cells is SR α . See Takebe et al., *Mol. Cell. Biol.* 8(1): 466-472 (1988). A suitable expression vector is pDSR α 2, which is further described below.

Enhancer Element

An enhancer sequence may be inserted into the vector to increase the transcription of a DNA sequence encoding a truncated GDNF protein of the present invention by higher eukaryotes. Enhancers are cis-acting elements of DNA, usually about from 10-300 bp in length, that act on the promoter to increase its transcription. Enhancers are relatively orientation and position independent. They have been found 5' and 3' to the transcription unit. Several enhancer sequences available from mammalian genes are known (e.g., globin, elastase, albumin, alpha-feto-protein and insulin). Typically, however, an enhancer from a virus will be used. The SV40 enhancer, the cytomegalovirus early promoter enhancer, the polyoma enhancer, and adenovirus enhancers are exemplary enhancing elements for the activation of eukaryotic promoters. While an enhancer may be spliced into the vector at a position 5' or 3' to truncated GDNF DNA, it is typically located at a site 5' from the promoter.

Transcription Termination

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and occasionally 3' untranslated regions of eukaryotic DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding truncated GDNF.

The construction of suitable vectors containing one or more of the above-listed components together with the desired truncated GDNF coding sequence is accomplished by standard ligation techniques. Isolated plasmids or DNA fragments are cleaved, tailored, and religated in the desired order to generate the plasmids required. To confirm that the correct sequences have been constructed, the ligation mixtures may be used to transform *E. coli*, and successful transformants may be selected by known techniques, such as ampicillin or tetracycline resistance as described above. Plasmids from the transformants are then prepared, analyzed by restriction endonuclease digestion, and/or sequenced to confirm the presence of the desired construct.

Vectors that provide for the transient expression of DNA encoding truncated GDNF in mammalian cells may also be used. In general, transient expression involves the use of an expression vector that is able to replicate efficiently in a host cell, such that the host cell accumulates many copies of the expression vector and, in turn, synthesizes high levels of the desired protein encoded by the expression vector. Transient expression systems, comprising a suitable expression vector and a host cell, allow for the convenient positive identification of proteins encoded by cloned DNAs, as well as for the rapid screening of such proteins for desired biological or physiological properties. Thus, transient expression systems are particularly useful in identifying variants of the protein.

Selection and Transformation of Host Cells

Host cells (e.g., bacterial, mammalian, insect, yeast, or plant cells) transformed with nucleic acid sequences for use in expressing a recombinant truncated GDNF protein are also provided by the present invention. The transformed host cell is cultured under appropriate conditions permitting the

expression of the nucleic acid sequence. The selection of suitable host cells and methods for transformation, culture, amplification, screening and product production and purification are well known in the art. See for example, Gething and Sambrook, *Nature* 293: 620-625 (1981), or alternatively, Kaufman et al., *Mol. Cell. Biol.*, 5 (7): 1750-1759 (1985) or Howley et al., U.S. Pat. No. 4,419,446. Truncated GDNF may be expressed in *E. coli* in accordance with the description of Lin et al. (U.S. Patent Application No. 07/855,413; Application No. PCT/US92/07888; WO 93/06116) which involved the expression of mature GDNF. Other exemplary materials and methods are discussed in further detail below. The transformed host cell is cultured in a suitable medium, and the expressed factor is then optionally recovered, isolated and purified from the culture medium (or from the cell, if expressed intracellularly) by an appropriate means known to those skilled in the art.

Suitable host cells for cloning or expressing the vectors herein are the prokaryote, yeast, or higher eukaryote cells as described above. Prokaryotic host cells include, but are not limited to, eubacteria, such as Gram-negative or Gram-positive organisms, for example, *E. coli*, *Bacilli* such as *B. subtilis*, *Pseudomonas* species such as *P. aeruginosa*, *Salmonella typhimurium*, or *Serratia marcescans*. Alternatively, *in vitro* methods of cloning, e.g., PCR or other nucleic acid polymerase reactions, are suitable.

In addition to prokaryotic host cells, eukaryotic microbes such as filamentous fungi or yeast may be suitable hosts for the expression of truncated GDNF proteins. *Saccharomyces cerevisiae*, or common baker's yeast, is the most commonly used among lower eukaryotic host microorganisms, but a number of other genera, species, and strains are well known and commonly available.

Suitable host cells for the expression of glycosylated truncated GDNF protein are derived from multicellular organisms. Such host cells are capable of complex processing and glycosylation activities. In principle, any higher eukaryotic cell culture might be used, whether such culture involves vertebrate or invertebrate cells, including plant and insect cells. Vertebrate cells are generally used as the propagation of vertebrate cells in culture (tissue culture) is a well known procedure. Examples of useful mammalian host cell lines include, but are not limited to, monkey kidney CV1 line transformed by SV40 (COS-7), human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture), baby hamster kidney cells, and Chinese hamster ovary cells. Other suitable mammalian cell lines include but are not limited to, HeLa,

mouse L-929 cells, 3T3 lines derived from Swiss, Balb-c or NIH mice, BHK or HaK hamster cell lines.

Similarly useful as host cells suitable for the present invention are bacterial cells. For example, the various strains of *E. coli* (e.g., HB101, DH5 α , DH10, and MC1061) are well-known as host cells in the field of biotechnology. Various strains of *Streptomyces spp.* and the like may also be employed. Presently preferred host cells for producing truncated GDNF proteins are bacterial cells (e.g., *Escherichia coli*) and mammalian cells (such as Chinese hamster ovary cells, COS cells, etc.)

The host cells are transfected and preferably transformed with the above-described expression or cloning vectors and cultured in a conventional nutrient medium. The medium may be modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. Transfection and transformation are performed using standard techniques which are well known to those skilled in the art and which are selected as appropriate to the host cells involved. For example, for mammalian cells without cell walls, the calcium phosphate precipitation method may be used. Electroporation, micro injection and other known techniques may also be used.

Culturing the Host Cells

Transformed cells used to produce truncated GDNF proteins of the present invention are cultured in suitable media. The media may be supplemented as necessary with hormones and/or other growth factors (such as insulin, transferrin, or epidermal growth factor), salts (such as sodium chloride, calcium, magnesium, and phosphate), buffers (such as HEPES), nucleosides (such as adenosine and thymidine), antibiotics (such as gentamicin), trace elements (defined as inorganic compounds usually present at final concentrations in the micromolar range), and glucose or other energy source. Other supplements may also be included, at appropriate concentrations, as will be appreciated by those skilled in the art. Suitable culture conditions, such as temperature, pH, and the like, are also well known to those skilled in the art for use with the selected host cells.

It is also possible that truncated GDNF proteins may be produced by homologous recombination, or with recombinant production methods utilizing control elements introduced into cells already containing DNA encoding GDNF. Homologous recombination is a technique originally developed for targeting genes

to induce or correct mutations in transcriptionally active genes (Kucherlapati, *Prog. in Nucl. Acid Res. and Mol. Biol.* 36:301 (1989)). The basic technique was developed as a method for introducing specific mutations into specific regions of the mammalian genome (Thomas et al., *Cell.* 44:419-428, 1986; Thomas and Capecchi, *Cell.* 51:503-512, 1987; Doetschman et al., *Proc. Natl. Acad. Sci.* 85:8583-8587, 1988) or to correct specific mutations within defective genes (Doetschman et al., *Nature.* 330:576-578, 1987). Exemplary homologous recombination techniques are described in U.S. 5,272,071 (EP 91 90 3051, EP Publication No. 505 500; PCT/US90/07642, International Publication No. WO 91/09955) the disclosure of which is hereby incorporated by reference.

Through homologous recombination, the DNA sequence to be inserted into the genome can be directed to a specific region of the gene of interest by attaching it to targeting DNA. The targeting DNA is DNA that is complementary (homologous) to a region of the genomic DNA. Small pieces of targeting DNA that are complementary to a specific region of the genome are put in contact with the parental strand during the DNA replication process. It is a general property of DNA that has been inserted into a cell to hybridize and therefore recombine with other pieces of endogenous DNA through shared homologous regions. If this complementary strand is attached to an oligonucleotide that contains a mutation or a different sequence of DNA, it too is incorporated into the newly synthesized strand as a result of the recombination. As a result of the proofreading function, it is possible for the new sequence of DNA to serve as the template. Thus, the transferred DNA is incorporated into the genome.

If the sequence of a particular gene is known, such as the nucleic acid sequence of GDNF, the pre-pro sequence or expression control sequence, a piece of DNA that is complementary to a selected region of the gene can be synthesized or otherwise obtained, such as by appropriate restriction of the native DNA at specific recognition sites bounding the region of interest. This piece serves as a targeting sequence upon insertion into the cell and will hybridize to its homologous region within the genome. If this hybridization occurs during DNA replication, this piece of DNA, and any additional sequence attached thereto, will act as an Okazaki fragment and will be backstitched into the newly synthesized daughter strand of DNA.

In the present invention, attached to these pieces of targeting DNA are regions of DNA which may interact with the expression of a GDNF protein. For

example, a promoter/enhancer element, a suppresser, or an exogenous transcription modulatory element is inserted in the genome of the intended host cell in proximity and orientation sufficient to influence the transcription of DNA encoding the desired truncated GDNF. The control element does not encode truncated GDNF, but instead controls a portion of the DNA present in the host cell genome. Thus, the expression of truncated GDNF proteins may be achieved not by transfection of DNA that encodes the truncated GDNF gene itself, but rather by the use of targeting DNA (containing regions of homology with the endogenous gene of interest) coupled with DNA regulatory segments that provide the endogenous gene sequence with recognizable signals for transcription of a truncated GDNF protein.

In accordance with the present invention, homologous recombination methods may also be used to modify a cell that contains a normally transcriptionally silent GDNF gene to produce a cell which expresses GDNF. The GDNF protein may then be processed to form a truncated GDNF protein(s).

Truncated GDNF Pharmaceutical Compositions

Truncated GDNF protein product pharmaceutical compositions typically include a therapeutically effective amount of a truncated GDNF protein product in admixture with one or more pharmaceutically and physiologically acceptable formulation materials. Suitable formulation materials include, but are not limited to, antioxidants, preservatives, coloring, flavoring and diluting agents, emulsifying agents, suspending agents, solvents, fillers, bulking agents, buffers, delivery vehicles, diluents, excipients and/or pharmaceutical adjuvants. For example, a suitable vehicle may be water for injection, physiological saline solution, or artificial cerebrospinal fluid (CSF), possibly supplemented with other materials common in compositions for parenteral administration. Neutral buffered saline or saline mixed with serum albumin are further exemplary vehicles.

The primary solvent in a vehicle may be either aqueous or non-aqueous in nature. In addition, the vehicle may contain other pharmaceutically-acceptable excipients for modifying or maintaining the pH, osmolarity, viscosity, clarity, color, sterility, stability, rate of dissolution, or odor of the formulation. Similarly, the vehicle may contain still other pharmaceutically-acceptable excipients for modifying or maintaining the stability, rate of dissolution, or rate of release of truncated GDNF protein product, or for promoting the absorption or penetration of truncated GDNF protein product

across the blood-brain barrier. Such excipients are those substances usually and customarily employed to formulate dosages for parenteral administration in either unit dose or multi-dose form or for direct infusion into the CSF by continuous or periodic infusion from an implanted pump.

5 Once the therapeutic composition has been formulated, it may be stored in sterile vials as a solution, suspension, gel, emulsion, solid, or dehydrated or lyophilized powder. Such formulations may be stored either in a ready to use form or in a form, e.g., lyophilized, requiring reconstitution prior to administration.

10 The optimal pharmaceutical formulation will be determined by one skilled in the art depending upon the route of administration and desired dosage. See for example, *Remington's Pharmaceutical Sciences*, 18th Ed. (1990, Mack Publishing Co., Easton, PA 18042) pages 1435-1712 the disclosure of which is hereby incorporated by reference. The composition may also involve particulate
15 preparations of polymeric compounds such as polylactic acid, polyglycolic acid, etc. or into liposomes. Hylauronic acid may also be used, and this may have the effect of promoting sustained duration in the circulation. Such compositions may influence the physical state, stability, rate of *in vivo* release, and rate of *in vivo* clearance of the present proteins and derivatives.

20 Other effective administration forms, such as parenteral slow-release formulations, inhalant mists, orally active formulations, or suppositories, are also envisioned. Current truncated GDNF protein product pharmaceutical compositions are formulated for parenteral administration, e.g., intracerebroventricular injection. Such parenterally administered therapeutic
25 compositions are typically in the form of a pyrogen-free, parenterally acceptable aqueous solution comprising truncated GDNF protein product in a pharmaceutically acceptable vehicle. One preferred vehicle is physiological saline.

30 It is also contemplated that certain formulations containing truncated GDNF protein product are to be administered orally. Truncated GDNF protein product which is administered in this fashion may be encapsulated and may be formulated with or without those carriers customarily used in the compounding of solid dosage forms. The capsule may be designed to release the active portion of the formulation at the point in the gastrointestinal tract when bioavailability is
35 maximized and pre-systemic degradation is minimized. Additional excipients may be included to facilitate absorption of truncated GDNF protein product.

Diluents, flavorings, low melting point waxes, vegetable oils, lubricants, suspending agents, tablet disintegrating agents, and binders may also be employed.

5 Administration of Truncated GDNF Protein Product

The truncated GDNF protein product may be administered parenterally via a subcutaneous, intramuscular, intravenous, transpulmonary, transdermal, intrathecal or intracerebral route. Protein growth factors that do not cross the blood-brain barrier may be given directly intracerebrally or otherwise in association with other elements that will transport them across the barrier. It is preferred that the truncated GDNF protein product is administered intracerebroventricularly or into the brain or spinal cord subarachnoid space. Truncated GDNF protein product may also be administered intracerebrally directly into the brain parenchyma. Slow-releasing implants in the brain containing the neurotrophic factor embedded in a biodegradable polymer matrix can also deliver truncated GDNF protein product. Truncated GDNF protein product may be administered extracerebrally in a form that has been modified chemically or packaged so that it passes the blood-brain barrier, or it may be administered in connection with one or more agents capable of promoting penetration of truncated GDNF protein product across the barrier. For example, a conjugate of NGF and monoclonal anti-transferrin receptor antibodies has been shown to be transported to the brain via binding to transferrin receptors. To achieve the desired dose of truncated GDNF protein product, repeated daily or less frequent injections may be administered, or truncated GDNF protein product may be infused continuously or periodically from a constant- or programmable-flow implanted pump. The frequency of dosing will depend on the pharmacokinetic parameters of the truncated GDNF protein product as formulated, and the route of administration.

Regardless of the manner of administration, the specific dose is typically calculated according to body weight or body surface area. For diseases involving the brain, the specific dose is typically calculated according to the approximate brain weight of the patient, which also may be estimated based on body weight or body surface area. Further refinement of the calculations necessary to determine the appropriate dosage for treatment involving each of the above mentioned formulations is routinely made by those of ordinary skill in the art, especially in light of the dosage information and assays disclosed herein.

Appropriate dosages may be ascertained through use of the established assays for determining dosages utilized in conjunction with appropriate dose-response data. The final dosage regimen involved in a method of treating a specific condition will be determined by the attending physician, considering various factors which

5 modify the action of drugs, e.g., the age, condition, body weight, sex and diet of the patient, the severity of any infection, time of administration and other clinical factors.

Truncated GDNF protein product of the present invention may also be employed, alone or in combination with other growth factors in the treatment of

10 nerve disease. For example, truncated GDNF protein product may be used in treating some forms of nerve disease in combination with nerve growth factor. In addition, other factors or other molecules, including chemical compositions, may be employed together with truncated GDNF protein product. In the treatment of Parkinson's Disease, it is contemplated that truncated GDNF protein product

15 be used by itself or in conjunction with the administration of Levodopa, wherein the truncated GDNF would enhance the production of endogenous dopamine and the neuronal uptake of the increased concentration of dopamine.

As stated above, it is also contemplated that additional neurotrophic or neuron nurturing factors will be useful or necessary to treat some neuronal cell

20 populations or some types of injury or disease. Other factors that may be used in conjunction with truncated GDNF include, but are not limited to: mitogens such as insulin, insulin-like growth factors, epidermal growth factor, vasoactive growth factor, pituitary adenylate cyclase activating polypeptide, interferon and somatostatin; neurotrophic factors such as brain derived neurotrophic factor,

25 neurotrophin-3, neurotrophin-4/5, neurotrophin-6, insulin-like growth factor, ciliary neurotrophic factor, acidic and basic fibroblast growth factors, fibroblast growth factor-5, transforming growth factor- β , cocaine-amphetamine regulated transcript (CART) and mature GDNF; and other growth factors such as epidermal growth factor, leukemia inhibitory factor,

30 interleukins, interferons, and colony stimulating factors; as well as molecules and materials which are the functional equivalents to these factors.

It is envisioned that the continuous administration or sustained delivery of a truncated GDNF protein product may be advantageous for a given treatment. While continuous administration may be accomplished via a mechanical means,

35 such as with an infusion pump, it is contemplated that other modes of continuous or near continuous administration may be practiced. For example, chemical

derivatization may result in sustained release forms of the protein which have the effect of continuous presence in the blood stream, in predictable amounts, based on a determined dosage regimen. Thus, truncated GDNF protein products include truncated GDNF protein derivatized to effectuate such continuous administration.

Truncated GDNF protein cell therapy, e.g., intracerebral implantation of cells producing truncated GDNF protein, is also contemplated. This embodiment of the present invention may include implanting into patients cells which are capable of synthesizing and secreting a biologically active form of truncated GDNF protein. Such truncated GDNF protein producing-cells may be cells which do not normally produce a neurotrophic factor but have been modified to produce truncated GDNF, or they could be cells whose ability to produce GDNF has been augmented by transformation with a polynucleotide suitable for the expression and secretion of truncated GDNF protein. In order to minimize a potential immunological reaction in patients from administering GDNF of a foreign species, it is preferred that the cells be of human origin and produce truncated human GDNF protein.

Implanted cells may be encapsulated to avoid infiltration of the cells into brain tissue. Human or non-human animal cells may be implanted in patients in biocompatible, semi-permeable polymeric enclosures or membranes to allow release of a truncated GDNF protein product, but that prevent destruction of the cells by the patient's immune system or by other detrimental factors from the surrounding tissue. Alternatively, the patient's own cells, transformed *ex vivo* to produce truncated GDNF, could be implanted directly into the patient without such encapsulation.

The methodology for the membrane encapsulation of living cells is familiar to those of ordinary skill in the art, and the preparation of the encapsulated cells and their implantation in patients may be accomplished. See for example, U.S. Patent Numbers 4,892,538; 5,011,472; and 5,106,627, the disclosures of which are hereby incorporated by reference. A system for encapsulating living cells is also described in PCT Application WO 91/10425 of Aebischer et al., specifically incorporated herein by reference. See also, PCT Application WO 91/10470 of Aebischer et al.; Winn et al., *Exper. Neurol.*, 113:322-329, 1991; Aebischer et al., *Exper. Neurol.*, 111:269-275, 1991; Tresco et al., *ASAIO*, 38:17-23, 1992, the disclosures of which are hereby incorporated by reference.

Truncated GDNF protein gene therapy *in vivo* is also envisioned, wherein a nucleic acid sequence encoding a truncated GDNF protein is introduced directly into the patient. For example, a nucleic acid sequence encoding a truncated GDNF protein is introduced into target cells via local injection of a nucleic acid construct with or without an appropriate delivery vector, such as an adeno-associated virus vector. Alternative viral vectors include, but are not limited to, retrovirus, adenovirus, herpes simplex virus and papilloma virus vectors. Physical transfer may be achieved *in vivo* by local injection of the desired nucleic acid construct or other appropriate delivery vector containing the desired nucleic acid sequence, liposome-mediated transfer, direct injection (naked DNA), receptor-mediated transfer (ligand-DNA complex), or microparticle bombardment (gene gun).

It should be noted that the truncated GDNF protein product formulations described herein may be used for veterinary as well as human applications and that the term "patient" should not be construed in a limiting manner. In the case of veterinary applications, the dosage ranges should be the same as specified above.

As a means of further characterizing truncated GDNF proteins of the present invention, antibodies can be developed which bind to the truncated GDNF protein such as to epitopes within the X-[Cys⁴¹-Cys¹³³]-Y amino acid sequence. One of ordinary skill in the art can use well-known, published procedures to obtain monoclonal and polyclonal antibodies, or recombinant antibodies, which specifically recognize and bind to the various proteins encoded by the amino acid sequences of the present invention. Such antibodies may then be used to purify and characterize truncated GDNF protein. Alternatively, the antibodies may be used as therapeutical inhibitors of the proteins to which they are directed.

Other aspects and advantages of the present invention will be understood upon consideration of the following illustrative examples. Example 1 addresses the expression of mature GDNF in a mammalian cell system and the preparation of truncated GDNF protein. Example 2 addresses the expression of mature GDNF in a bacterial cell system. Example 3 addresses the expression of various truncated GDNF proteins in a bacterial cell system. Example 4 compares the biological activity of the mature GDNF protein and truncated GDNF protein in an assay for dopaminergic neuron neurotrophic activity.

EXAMPLES

5

Example 1

Expression of Mature Human GDNF in CHO Cells
and the Purification of CHO cell-derived Truncated GDNF Protein

Materials

10 The following materials are used in the expression of human GDNF in dihydrofolate reductase-deficient CHO cells (CHO⁻ cells; for example, as described by Urlaub and Chasin, *Proc. Natl. Acad. Sci., USA* 77(7): 4216-4220 (1980)).

CHO⁻ medium contained: Dulbecco's Modified Eagle's Medium (DMEM) -
15 high glucose (Gibco/BRL); 5% fetal bovine serum (HyClone); MEM non-essential amino acids (1%) (Gibco/BRL); hypoxanthine/thymidine (1%) (Gibco/BRL); and glutamine/penicillin/streptomycin (1%) (Irvine Scientific).

Selective medium contained: DMEM (high glucose); 5% dialyzed fetal bovine serum (HyClone); MEM non-essential amino acids; and
20 glutamine/penicillin/streptomycin.

2X HEPES-buffered saline (HBS) contained: 280 mM NaCl; 10 mM KCl; 1.5 mM Na₂HPO₄; 12 mM dextrose; and 50 mM HEPES.

Tris-buffered saline plus Tween (TBST) contained: 137 mM NaCl; 20 mM Tris/HCl pH 7.5; and 0.1% Tween-20.

25

Methods

Transfection and Selection

CHO⁻ cells (passage 20) were seeded into 60mm tissue culture dishes (Falcon) at a density of 8×10^5 cells per dish in CHO⁻ growth medium. On the
30 following day, about three hours prior to transfection, the medium on the cells was replaced with fresh medium.

Plasmid constructs containing the appropriate GDNF cDNA were prepared using well known techniques. For example, the plasmid construct pDSRα2 was prepared substantially in accordance with the process described in the co-owned
35 and copending U. S. Patent Application Serial Number 501,904 filed March 29, 1990 (also see, European Patent Application No. 90305433, Publication No.

EP 398 753, filed May 18, 1990 and WO 90/14363 (1990), the disclosures of which are hereby incorporated by reference. An exemplary plasmid map which illustrates the structural organization of the vector is depicted in Figure 2. It will be appreciated by those skilled in the art that a variety of nucleic acid sequences encoding the mature GDNF protein, such as the sequences depicted in Figures 1, 3 and 4, may also be used.

A HindIII-XbaI DNA fragment containing the human GDNF coding sequences and the consensus Kozak sequences, CCACC(ATG) was retrieved by restriction enzyme digestion from a pcDNA3 based expression vector (Invitrogen, San Diego, CA). The DNA fragment was directly cloned into HindIII/XbaI cut pDSR α 2. The resulting plasmid was called pSW5. The plasmid DNA of pSW5 was linearized at the PvuI site prior to transfection.

pDSR α 2 (Figure 2) is a derivative of the plasmid pCD (Okayama & Berg, *Mol. Cell Biol.* 3: 280-289, 1983) with three main modifications: (i) the SV40 polyadenylation signal has been replaced with the signal from the α -subunit of bovine follicular stimulating hormone, α -bFSH (Goodwin et al., *Nucleic Acids Res.* 11: 6873-6882, 1983); (ii) a mouse dihydrofolate reductase minigene (Gasser et al., *Proc. Natl. Acad. Sci.* 79: 6522-6526, 1982) has been inserted downstream from the expression cassette to allow selection and amplification of the transformants; and (iii) a 267 bp fragment containing the "R-element" and part of the "U5" sequences of the long terminal repeat (LTR) of human T-cell leukemia virus type I (HTLV-I) has been cloned and inserted between the SV40 promoter and the splice signals as described previously (Takebe et al., *Mol. Cell Biol.* 8: 466-472, 1988).

Solutions of DNA were prepared containing a final concentration of 3.0 μ g/dish GDNF-plasmid DNA, 7.0 μ g/dish mouse kidney genomic carrier DNA (Clontech), 25 μ l/dish 2.5M CaCl₂, and sterile distilled water to a final volume of 250 μ l/dish. DNA solutions containing pDSR α 2 vector DNA or carrier DNA alone were similarly prepared as positive and negative controls, respectively. The DNA solutions were added dropwise to an equal volume of 2X HEPES-buffered saline while passing air bubbles through the solution. The DNA/HBS solutions were incubated at room temperature for 30 minutes.

The medium was removed from the CHOD⁺ cell cultures, and 500 μ l of the DNA solutions were added per dish. The dishes were incubated at room temperature for 30 minutes, after which time CHOD⁺ medium (5.0 ml.) was added to each dish. The dishes were then incubated at 37°C overnight.

On the following day the medium was replaced with fresh CHO^d medium. The next day, when the cells had reached confluence, the cultures were trypsinized and replated in 100 mm dishes (Falcon) at a ratio of 1x60 mm dish to 8x100 mm dishes. Cells were replated in selective medium. The cultures were re-fed with fresh medium every two to three days.

After 15 days, colonies of transfected cells were isolated using glass cloning cylinders, trypsinized, and replated into 24-well dishes (Falcon). A total of 40 colonies was isolated from the GDNF/pSW5-transfected cells. The remaining cells on the dishes were trypsinized, pooled, and replated into two 100 mm dishes (one pool for each DNA construct).

Screening of transfected cells:

The 24-well and pool cultures were grown to confluence, at which time the growth medium was removed and replaced with serum-free medium (400 μ l/well or 4 ml/dish). Cells were incubated for 48 hours, and the conditioned medium was harvested. The conditioned medium samples were analyzed for GDNF protein expression by western blot. Aliquots of conditioned medium (20 μ l or 40 μ l) were diluted with electrophoresis sample buffer (with or without β -mercaptoethanol). Samples containing β -mercaptoethanol were boiled for three minutes (reducing conditions). Both reduced and non-reduced samples were run on 16% Tris-glycine gels (Novex). Gels were electroblotted onto nitrocellulose filters (Schleicher and Schuell BA-83, 0.2 μ). The blots were rinsed with TBST and then incubated in a blocking solution of 5% dried milk (Carnation) in TBST for 30 minutes at room temperature. Blots were then treated with GDNF antiserum (rabbit polyclonal antisera raised against *E. coli*-derived GDNF; 1:1000 in 5% milk/TBST) for one hour at room temperature. The blots were then rinsed with TBST and washed 1x10 minutes and 2x5 minutes with 1% milk/TBST. They were then treated with anti-rabbit Ig-horse radish peroxidase-conjugated secondary antibody (1:15,000 in 1% milk/TBST) for 20 minutes. Blots were rinsed and washed with TBST 1x20 minutes and 2x10 minutes, followed by treatment with ECL reagents (Amersham) for one minute and exposure to Hyperfilm-ECL (Amersham).

The following process describes the purification of CHO-expressed GDNF and a CHO-derived clipped GDNF homodimer, from one liter of conditioned media. Because of significant protease action in the CHO medium, clipping the chain at

residue 31, the procedure may include the use of a protease inhibitor during purification.

Step 1. Bead Chromatography:

5 Serum free conditioned media was made 20 mM 2-[N-Morpholino] ethane sulfonate (MES), pH 6.0, by adding one fiftieth volume of 1 M MES, pH 6.0. Twenty five milliliters of SP Sepharose Big Bead resin (Pharmacia), equilibrated with 20 mM MES, pH 6.0, was added and stirred at 4°C for one hour. The resin was collected by allowing it to settle and decanting off the conditioned
10 media. The decanted media was filtered through a fritted disc filter to recover any unsettled resin. The settled resin and that recovered by filtration was resuspended and poured into a 2.5 cm diameter column and washed with three column volumes of 0.15 M NaCl, 20 mM MES, pH 6.0 (A buffer). Protein was eluted with a 300 ml gradient from A buffer to 1.0 M NaCl, 20 mM MES, pH 6.0
15 (B buffer), at a flow rate of 0.2 column volumes/minute with absorbance monitored at 280 nm. Fractions containing 1.1 column volumes were collected. The presence of GDNF in the fractions was detected by Western blotting analysis. Fractions containing GDNF were pooled for further purification. GDNF eluted between 0.3 and 0.6 M NaCl.

Step 2. HPLC C4 Chromatography:

The pool from Step 1 was made 0.1% (v/v) trifluoroacetic acid (TFA), vacuum filtered through a 0.45 micron filter, and applied to a Vydac C4 column (0.46 X 25 cm) conditioned with aqueous 10% acetonitrile, 0.1% TFA (A
25 buffer). Protein was eluted with a 2%/minute linear gradient over 50 minutes from A buffer to aqueous 90% acetonitrile, 0.1% TFA (B buffer) with absorbance measured at 280 nm. One milliliter fractions were collected, and the presence of GDNF was detected by Western blotting analysis. GDNF was eluted between 45% and 55% acetonitrile. Fractions were taken to dryness in vacuum.

Step 3. High performance S Chromatography:

Fractions containing GDNF from Step 2 were resolubilized in one milliliter of 0.15 M NaCl, 10 mM Tris, pH 8.0, and applied to a 0.75 X 7.5 cm TSK-Gel SWP high performance S column (Toso Haas). A linear gradient of
35 0.4%/minute was run from 0.15 M NaCl, 10 mM Tris, pH 8.0 (A buffer) to 1.0 M NaCl, 10 mM Tris, pH 8.0 (B buffer) over 50 minutes at a flow rate of

1 ml/min. One minute fractions were collected with absorbance measured at 280 nm. At 35% B buffer, the gradient was changed to 6.5%/minute over 10 minutes. Western blot analysis of the fractions showed four major GDNF components. Three of the components eluted during the 0.4%/minute gradient and the fourth was eluted during the 6.5%/minute gradient. Appropriate pools were made of similar components and submitted for sequencing. Sequencing analyses identified an approximately 29-36 kD pool as [Arg³²-Ile¹³⁴] truncated GDNF protein. The component at approximately 38-40 kD was identified as an [Arg³²-Ile¹³⁴] truncated GDNF/mature GDNF heterodimer. Finally, the approximately 41-44 kD component which was isolated during the latter portion of the gradient was identified by sequencing as the mature GDNF homodimer.

Example 2

Mature Human GDNF Produced in *E. coli*

The bacterial expression of mature human GDNF may be achieved in accordance with the process described in Lin et al. (U.S. Patent Application No. 08/182,183 filed May 23, 1994 and its parent applications; PCT/US92/07888 filed September 17, 1992 (WO 93/06116); and European Patent Application No. 92921022.7 (Publication No. EP 610 254); the disclosures of which are hereby incorporated by reference. Based upon the description of the present invention, those of ordinary skill in the art will appreciate that a variety of materials and methods may readily be used or adapted for suitable expression in *E. coli* and other bacteria. For example, alternate polynucleotides, such as those depicted in Figures 1, 3 and 4, may be used in the expression process.

Refolding and Purification of Expressed Mature GDNF

The transformed cells were processed at 5°C (unless otherwise noted) as follows: cell paste (30 gm) was suspended into a final volume of 200 milliliters using 25 mM Tris, pH 8.5 containing 5 mM EDTA, to yield a final cell slurry of 15% (w/v). The cells were thoroughly dispersed using a Biospec hand-held low shear homogenizer. The slurry was passed twice through a microfluidizer at 14,500 psi to break the cells and release inclusion bodies. The resulting

homogenate was then centrifuged for 30 minutes at 16,000xg. The pellet of inclusion bodies resulting from the centrifugation was washed by resuspension in chilled water to a final volume of 240 milliliters using the Biospec homogenizer, as before, to form a slurry. A sample of this slurry was kept for

- 5 HPLC analysis of the GDNF expression level. The remaining slurry was centrifuged for 30 minutes at 16,000xg. The supernatant was discarded, and a small amount of cold water was added to the centrifuge bottle and gently swirled to remove the loosely formed membrane layer on top of the inclusion bodies pellet. The pellet was resuspended with the Biospec homogenizer using a
- 10 sufficient volume of cold water to yield a concentration of 2 mg/ml of GDNF. The inclusion bodies were then solubilized by mixing the final inclusion bodies suspension (25 ml) and 8M guanidine HCl (25 ml) containing 180 mM cysteine HCl, and 50 mM Tris HCl, pH 8.7. The solubilization mixture was stirred at 25°C for 60 to 90 minutes after which it was poured, with mixing, into 2 M
- 15 urea (450 ml, at 5°C) containing 20 mM Tris HCl, pH 8.75 and 0.2 M guanidine HCl. This refold mixture was slowly stirred at 5°C for 72 hours.

The refolded GDNF was partially purified as follows: 20 mM sodium acetate buffer (250 ml, pH 5) at 5°C was added with rapid stirring to the refold mixture, and the pH was adjusted to 5 with glacial acetic acid. The resulting

20 precipitate was removed by centrifugation at 13,600xg for 45 minutes at 5°C. The supernatant from this centrifugation was used as the load solution for the next purification step involving cation exchange chromatography with SP-big bead resin (Pharmacia). The column was operated at 5°C using 20 mM sodium acetate (pH 5) as the equilibration, rinsing, and elution buffer system. A bed of

25 resin (5 ml) was sanitized with 5 column volumes (CV) of 0.2N NaOH and then equilibrated with the acetate buffer (5 CV). The load solution (190 ml) was applied to the column at 0.5 CV/minute followed by a 10 CV rinse with acetate buffer at the same flow rate. The GDNF was then eluted off the resin with a 20 CV linear gradient of NaCl from 0.3 M to 0.9M in the acetate buffer at a flow rate of

30 0.1 CV/minute. The column eluate was monitored by absorbance at 280 nm and collected as fractions which were assayed by SDS-PAGE. The fractions containing GDNF were pooled from the front of the GDNF peak at 10% peak height through to the back of the peak to 10% peak height. The protein in this pool was entirely GDNF and, depending upon the production strain used, contained 32% to 12%

35 contamination as modified GDNF forms. The pool was then dialyzed against PBS or other formulation buffers and, in some cases, concentrated by ultrafiltration

to 25 mg/ml. Both wild type and analogue forms of GDNF purified by this procedure were characterized by reverse phase HPLC, cation exchange HPLC, mass spectrometry, and endotoxin levels in order to compare purities of the preparations in relation to the corresponding production strains.

5

Example 3

Recombinant Production of Truncated GDNF in *E. coli*

10 Exemplary truncated GDNF proteins were produced substantially in accordance with the techniques described in Lin et al. (U.S. Patent Application No. 08/182,183 filed May 23, 1994, *supra*). Alternative bacterial expression materials and methods, as described above, may also be used. The *E. coli*-expressed truncated GDNF proteins included the [Pro²³-Ile¹³⁴], [Arg³²-Ile¹³⁴], and [Gly³³-Ile¹³⁴] truncated GDNF proteins as depicted in Figures 5, 6 and 7, respectively. The polynucleotides encoding these exemplary truncated GDNF proteins were constructed as depicted in Figures 5, 6 and 7, but corresponding polynucleotides such as those depicted in Figures 1, 3 and 4 may also be used. The polynucleotides were constructed by standard PCR procedures as described in *PCR Technology, Principles and Applications for DNA Amplification*, Henry A. Erlich, ed., Stockton Press, NY, 1989 (Chapter 6, Using PCR to Engineer DNA) the disclosure of which is hereby incorporated by reference.

25

Example 4

Bioassay for Dopaminergic Neuron Neurotrophic Activity

The *E. coli*-expressed [Pro²³-Ile¹³⁴], [Arg³²-Ile¹³⁴], [Gly³³-Ile¹³⁴] and [Lys³⁷-Ile¹³⁴] truncated GDNF proteins of Example 3 and the CHO-derived [Arg³²-Ile¹³⁴] truncated GDNF protein of Example 1 were assessed on a qualitative basis for their ability to promote dopamine uptake by substantia nigra dopaminergic neurons.

35 Materials

The following materials are used in an assay to assess the survival of dopaminergic neurons in the presence of truncated GDNF proteins:

Cell Culture Media

High glucose Dulbecco's Modified Eagle's Medium (DMEM; catalog #11965-092), Ham's F12 medium (F12; #11765-021), Leibovitz's L15 medium without sodium bicarbonate (#41300-039), B27 medium supplement (#17504-010), penicillin/streptomycin (#15070-014), L-glutamine (#25030-016), Dulbecco's phosphate-buffered saline (D-PBS; #14190-052), Hank's balanced salt solution with calcium and magnesium salts (HBSS; #24020-026), N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid (HEPES; #15630-015), mouse laminin (#23017-015), and bovine serum albumin fraction V (#110-18-017) were all from GIBCO, Grand Island, NY. Heat-inactivated horse serum was from HyClone, Logan, Utah. Conalbumin (C-7786), poly-L-ornithine hydrobromide (P-3655), bovine insulin (I-5500), human transferrin (T-2252), putrescine (P-6024), progesterone (P-6149), sodium selenite (S-9133), metrizamide (M-3383) were all from Sigma Chemical Company, Saint-Louis, MO. Papain, deoxyribonuclease I (DNAase) and ovalbumin (Papain dissociation system) were from Worthington Biochemicals, Freehold, NJ.

Falcon sterile 96-well microplates (#3072), tissue culture plastic ware and polypropylene centrifuge tubes were from Becton-Dickinson, Oxnard, CA. Nunc Lab-Tek tissue culture chamber coverglasses (#136439) were from Baxter, Irvine, CA; 20 μ m (#460) nylon mesh was from Tetko, Elmsford, NY; and 4" dissecting forceps and 4" dissecting scissors were from Roboz Surgical, Washington, DC.

Antibodies and Related Reagents

Polyclonal rabbit anti-tyrosine hydroxylase antibodies (TE101) were from Eugene Tech, Ridgefield Park, NJ; polyclonal rabbit anti-neuronal-specific enolase antibodies (NSE, AB951) were from Chemicon, Temecula, CA; and biotinylated goat anti-rabbit IgG and peroxidase-conjugated avidin/biotin complex (ABC Elite; Vectastain kit PK-6100) were from Vector Laboratories, Burlingame, CA. 3',3'-diaminobenzidine was from Cappel Laboratories, West Chester, PA. Superblock blocking buffer in PBS (#37515) was from Pierce Chemical Company, Rockford, IL. Triton X-100 (X100), Nonidet P-40 (N6507) and hydrogen peroxide (30%, v/v; H1009) were from Sigma. GBR-12909 dopamine uptake inhibitor (D-052) was from RBI, Natick, MA. 3 H-

dopamine (tritiated dopamine, NE-131; 21 Ci/mmol) was from New England Nuclear, Boston, MA. Optiphase Supermix scintillation cocktail was from Wallac, Turku, Finland. White ViewPlate-96 microplates (#6005182) were from Packard Instruments Corporation, Meriden, CT. All other reagents were obtained from Sigma Chemical Company, unless otherwise specified.

Preparation of Media

The basal medium was prepared as a 1:1 mixture of DMEM and F12 medium, and was supplemented with B27 medium supplement added as a 50-fold concentrated stock solution. L-glutamine was added at a final concentration of about 2 mM, penicillin at about 100 IU/l, and streptomycin at about 100 mg/l. Heat-inactivated horse serum was added to a final concentration of about 15 percent. After mixing, the pH was adjusted to about 7.3, and the media were kept at 4°C. The media were prepared fresh just before use in order to minimize inter-experimental variations. Plastic pipettes and containers were used throughout to minimize protein adsorption.

Culture Substratum

To encourage optimal attachment of substratum neurons and neurite outgrowth, microtiter plate surfaces (the culture substratum) were modified by sequential coatings with poly-L-ornithine and laminin, as follows. The plate surfaces were completely covered with a 0.1 mg/ml sterile solution of poly-L-ornithine in 0.1 M boric acid (pH 8.4) for at least one hour at room temperature, followed by a sterile wash with Super-Q water. The water wash was then aspirated, and a 1.0 µg/ml solution of mouse laminin in PBS was added and incubated at 37°C for two hours. These procedures were conducted just before using the plates in order to ensure reproducibility of the results.

Preparation of Embryonic Rat Substantia Nigra Cultures

Embryonic rat brains were used as the source of substantia nigra dopaminergic neurons. Timed-pregnant Sprague-Dawley rats at embryonic day 15 were used. A maximum of 36 embryos (about three litters) were processed per experiment. The pregnant rats were killed by exposure to CO₂, their abdominal cavities opened with dissecting scissors, and the fetuses were removed from the uteri. Fetal brains were then dissected and cleaned of blood and meninges, and the ventral tegmental area containing the substantia nigra was

dissected using well-defined anatomical landmarks (Altman and Bayer, *Atlas of Prenatal Rat Brain Development*, CRC Press, Boca Raton, FL, 1995). The tissues were collected in ice-cold D-PBS, transferred into 10 milliliters of dissociation medium (120 units papain and 2000 units DNAase in HBSS) and then incubated for 45 minutes at about 37°C on a rotary platform shaker at about 200 rpm. The cells were then dispersed by trituration through fire-polished Pasteur pipettes, sieved through a 20 µm Nitex mesh to discard undissociated tissue, and centrifuged for five minutes at 200xg using an IEC clinical centrifuge. The resulting cell pellet was resuspended into HBSS containing ovalbumin and about 500 units DNAase, layered on top of a 4% ovalbumin solution (in HBSS) and centrifuged for about 10 minutes at 500xg. The final pellet was resuspended in complete culture medium (see above), adjusted to about 28,000 cells/ml, and seeded in aliquots (90 µl) into the 6 mm-wells of the 96-well microplates previously coated with polyornithine and laminin. Attachment of cells occurred rapidly, and the plating efficiency was about 75 percent.

Immunohistochemistry of Dopaminergic Neurons

An indirect immunoperoxidase method described by Louis et al. (*J. Pharmacol. Exp. Therap.*, 262:1274-1283, 1992; *Science*, 259:689-692, 1993) was used with slight modifications, as follows, to characterize the dopaminergic neurons in cultures of substantia nigra. Cultures were fixed for about 30 minutes at room temperature with 4% paraformaldehyde in D-PBS, pH 7.4, followed by three washes in D-PBS (200 µl per 6-mm well). The fixed cultures were then incubated in Superblock blocking buffer in PBS, containing 1% NP-40 to increase the penetration of the antibodies. The primary rabbit anti-tyrosine hydroxylase antibodies were then applied at a dilution of about 1:2000 in the same buffer and incubated for one hour at 37°C on a rotary shaker. After three washes with D-PBS, the bound antibodies were detected using goat-anti-rabbit biotinylated IgG at about a 1:500 dilution; these secondary antibodies were incubated with the cells for about one hour at 37°C. The cells were then washed three times with D-PBS, and the secondary antibodies were detected with avidin-biotin-peroxidase complex diluted at 1:500, and the cells were incubated for about 45 minutes at 37°C. After three more washes with D-PBS, the cultures were reacted for 5-20 minutes in a

solution of 0.1 M Tris-HCl, pH 7.4, containing 0.04% 3',3'-diaminobenzidine-(HCl)₄, 0.06 percent NiCl₂ and 0.02 percent hydrogen peroxide.

Determining Neuronal Survival

- 5 Substantia nigra cultures were fixed and processed for immunostaining as described above, and then examined with bright-light optics at 200X magnification. The number of neurons stained for tyrosine hydroxylase was counted in the entire 6-mm well of the 96-well microplates. Viable neurons were characterized as having a regularly-shaped cell body, with a major axon-like process and several dendrite-like processes. Neurons showing signs of
- 10 degeneration, such as having irregular, vacuolated perikarya or fragmented neurites, were excluded from the counts (most of the degenerating neurons, however, detached from the culture substratum). Dopaminergic neuron cell numbers were expressed either as TH-positive neurons/6-mm well or as the
- 15 fold-change relative to control dopaminergic neuron density.

Determining Dopamine Uptake

- Dopamine uptake was determined in cultures of 15-day-old embryonic rat substantia nigra neurons that had been established in white ViewPlate-96
- 20 microplates. The cultures were washed with pre-warmed uptake buffer (about 100 μ l) which consists of a modified Krebs-Ringer solution, pH 7.4, containing about 120 mM NaCl, 4.7 mM KCl, 1.8 mM CaCl₂, 1.2 mM MgSO₄, 32 mM NaHPO₄, 1.3 mM EDTA, and 5.6 mM D-glucose. The uptake buffer also contained 1 mM ascorbic acid and 50 μ M pargyline to prevent the oxidation of dopamine.
- 25 The cells were then preincubated at 37°C for about 10 minutes in uptake buffer. Tritiated dopamine (³H-DA, 21 Ci/mmol) was then added to the substantia nigra cultures at a concentration of about 50 nM in 75 μ l of uptake buffer, and the cultures were incubated for about 60 minutes at 37°C. Non-specific dopamine uptake was determined by incubating the cultures with uptake buffer containing
- 30 the dopamine uptake inhibitor GBR-12909 (1 μ M). Non-specific uptake represented less than about one percent of total uptake. The uptake assays were arrested by aspiration of the incubation medium followed by three rapid washes with ice-cold uptake buffer (about 120 μ l). The cells were then lysed by addition of Optiphase Supermix scintillation cocktail (200 μ l), and radioactivity
- 35 was determined by scintillation spectrometry using a Wallac MicrobetaPlus 96-well microplate counter (i.e., dopamine uptake is analyzed by scintillation

counting of the retained tritium in the cultures). The results are expressed either as dpm/6-mm well or as the fold-change relative to control cultures.

Assays

5 Dopaminergic Neuron Survival and Morphological Development

Cultures of embryonic day 15 (E15) rat substantia nigra enriched in dopaminergic neurons were used to demonstrate the effect of truncated GDNF proteins on the survival of dopaminergic neurons. The cultures were grown in polyornithine- and laminin-coated 96-well microplates for up to six days alone or in the presence of various concentrations (ranging from about 1 pg/ml to about 10 ng/ml) of the following proteins: *E. coli*-expressed mature hGDNF; *E. coli*-expressed [Pro²³-Ile¹³⁴], [Arg³²-Ile¹³⁴], [Gly³³-Ile¹³⁴] and [Lys³⁷-Ile¹³⁴] truncated GDNF proteins; CHO cell-expressed mature hGDNF; and CHO cell-derived [Arg³²-Ile¹³⁴] truncated GDNF protein. The culture medium consisted of DMEM/F12 supplemented with 15% heat-inactivated horse serum (E15 cultures) or 2.5% heat-inactivated horse serum, D-glucose, HEPES, insulin and transferrin (P6 cultures). Immunostaining for tyrosine hydroxylase (TH), the rate-limiting enzyme in dopamine biosynthesis, was used as a marker for dopaminergic neurons. Since noradrenergic neurons in the rhombencephalon also stain positive for TH, great care was taken to dissect an area restricted to the ventral tegmentum of the mesencephalon and to avoid the more caudal regions containing the noradrenergic cell bodies. After six days, the E15 cultures typically consisted of about 70% neurons as identified by neuronal specific enolase immunostaining (described above) and 30% non-neuronal cells (which had a flattened, phase-dark appearance); dopaminergic neurons represented about 10-15% of the neuron population.

After six days, the cultures were fixed with paraformaldehyde and immunostained for tyrosine hydroxylase, a marker that identifies dopaminergic neurons in these cultures. All the tyrosine-hydroxylase-positive neurons present in a 6-mm well were counted under brightfield optics. Three to six different wells were analyzed for each experimental condition. The results were expressed as the percentages of the number of tyrosine-hydroxylase-positive neurons found in control cultures.

Cultures of E15 substantia nigra treated with 1.0 ng/ml of GDNF, CHO cell-expressed GDNF or *E. coli*-expressed GDNF, contained about 38% and 27% more TH-immunoreactive neurons, respectively, than untreated control

cultures, indicating that both GDNF species promote the survival of dopaminergic neurons. Cultures of E15 substantia nigra treated with 1.0 ng/ml of truncated GDNF protein showed a similar increase in the number of TH-positive neurons in cultures after six days *in vitro* : 42% for CHO cell-derived [Arg³²-Ile¹³⁴] truncated GDNF protein; and 26% and 17% for *E. coli*-expressed [Arg³²-Ile¹³⁴] and [Gly³³-Ile¹³⁴] truncated GDNF proteins, respectively.

Comparison of control cultures and cultures treated with mature and truncated GDNF proteins also revealed pronounced effects of all the GDNF proteins on the morphological differentiation of dopaminergic neurons. The effects of the [Arg³²-Ile¹³⁴] and [Gly³³-Ile¹³⁴] truncated GDNF proteins were identical to their respective mature GDNF protein counterparts. TH-immunoreactive neurons in all GDNF-treated cultures possessed significantly more complex and extensive neuritic arborization, as well as a higher degree of neurite branching and an overall larger soma size, than did TH-positive neurons in control cultures.

Dopamine Uptake

Dopamine uptake measures the number and activity of high affinity dopamine reuptake transporter sites and reflects the functional differentiation of dopaminergic neurons. Dopamine uptake was measured in cultures of E15 rat substantia nigra after six days *in vitro* either with or without mature GDNF or truncated GDNF proteins. In these cultures, dopamine uptake had the pharmacological profile characteristic of dopaminergic neurons, i.e., it was nearly completely blocked (more than 98 percent) by 1.0 μ M GBR-12909, a dopamine transporter inhibitor specific for dopaminergic neurons (ID₅₀ = 20 nM). This indicates that dopamine uptake measurements do not reflect the presence of contaminating noradrenergic neurons, which can take up dopamine through norepinephrine transporters but are not sensitive to GBR-12909 inhibition. The effects of CHO-cell-expressed mature GDNF and the CHO-derived [Arg³²-Ile¹³⁴] truncated GDNF protein were identical: about 65% increase, with an ED₅₀ of about 20 pg/ml. *E. coli*-expressed [Pro²³-Lys³⁷ Δ Asn³⁷-Ile¹³⁴] truncated GDNF protein, as depicted in Figure 5, demonstrated a 65% increase, with an ED₅₀ of about 40 pg/ml. The effects on dopamine uptake of the *E. coli*-expressed mature protein and the *E. coli*-expressed [Arg³²-Ile¹³⁴], [Gly³³-Ile¹³⁴] and [Lys³⁷-Ile¹³⁴] truncated GDNF proteins were the same: about 50% increases, with ED₅₀s of about 50 pg/ml.

These results indicate that the truncated GDNF proteins act as potent survival-promoting and differentiation-inducing factors for substantia nigra dopaminergic neurons. As such, they are envisioned to be particularly useful in the treatment of Parkinson's disease, a neurological disorder characterized by decreased emotional acuity, slowing of both voluntary and involuntary muscle movement, muscular rigidity, and tremor. Such symptoms are attributed to the progressive degeneration of dopamine-producing neurons located in the substantia nigra. Degeneration of these neurons ("dopaminergic neurons") results in a decrease of dopamine in an adjacent region of the brain called the striatum.

10

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Hu, Sylvia
- (ii) TITLE OF INVENTION: Truncated Glial Cell Line-Derived Neurotrophic Factor
- (iii) NUMBER OF SEQUENCES: 50
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: AMGEN INC.
 - (B) STREET: 1840 DeHavilland Drive
 - (C) CITY: Thousand Oaks
 - (D) STATE: California
 - (E) COUNTRY: United States of America
 - (F) ZIP: 91320
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Curry, Daniel R.
 - (B) REGISTRATION NUMBER: 32,727
 - (C) REFERENCE/DOCKET NUMBER: A-357
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 805-447-8102
 - (B) TELEFAX: 805-499-8011
 - (C) TELEX:

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 402 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (ix) FEATURE:
 - (A) NAME/KEY: CDS
 - (B) LOCATION: 1..402

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

TCA	CCA	GAT	AAA	CAA	ATG	GCA	GTG	CTT	CCT	AGA	AGA	GAG	CGG	AAT	CGG	48
Ser	Pro	Asp	Lys	Gln	Met	Ala	Val	Leu	Pro	Arg	Arg	Glu	Arg	Asn	Arg	
1				5					10					15		
CAG	GCT	GCA	GCT	GCC	AAC	CCA	GAG	AAT	TCC	AGA	GGA	AAA	GGT	CGG	AGA	96
Gln	Ala	Ala	Ala	Ala	Asn	Pro	Glu	Asn	Ser	Arg	Gly	Lys	Gly	Arg	Arg	
			20					25				30				
GGC	CAG	AGG	GGC	AAA	AAC	CGG	GGT	TGT	GTC	TTA	ACT	GCA	ATA	CAT	TTA	144
Gly	Gln	Arg	Gly	Lys	Asn	Arg	Gly	Cys	Val	Leu	Thr	Ala	Ile	His	Leu	
			35				40					45				
AAT	GTC	ACT	GAC	TTG	GGT	CTG	GGC	TAT	GAA	ACC	AAG	GAG	GAA	CTG	ATT	192
Asn	Val	Thr	Asp	Leu	Gly	Leu	Gly	Tyr	Glu	Thr	Lys	Glu	Glu	Leu	Ile	
			50			55					60					
TTT	AGG	TAC	TGC	AGC	GGC	TCT	TGC	GAT	GCA	GCT	GAG	ACA	ACG	TAC	GAC	240
Phe	Arg	Tyr	Cys	Ser	Gly	Ser	Cys	Asp	Ala	Ala	Glu	Thr	Thr	Tyr	Asp	
					70				75					80		
AAA	ATA	TTG	AAA	AAC	TTA	TCC	AGA	AAT	AGA	AGG	CTG	GTG	AGT	GAC	AAA	288
Lys	Ile	Leu	Lys	Asn	Leu	Ser	Arg	Asn	Arg	Arg	Leu	Val	Ser	Asp	Lys	
				85				90						95		
GTA	GGG	CAG	GCA	TGT	TGC	AGA	CCC	ATC	GCC	TTT	GAT	GAT	GAC	CTG	TCG	336
Val	Gly	Gln	Ala	Cys	Cys	Arg	Pro	Ile	Ala	Phe	Asp	Asp	Asp	Leu	Ser	
			100				105						110			
TTT	TTA	GAT	GAT	AAC	CTG	GTT	TAC	CAT	ATT	CTA	AGA	AAG	CAT	TCC	GCT	384
Phe	Leu	Asp	Asp	Asn	Leu	Val	Tyr	His	Ile	Leu	Arg	Lys	His	Ser	Ala	
			115			120						125				
AAA	AGG	TGT	GGA	TGT	ATC											402
Lys	Arg	Cys	Gly	Cys	Ile											
			130													

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 134 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Ser	Pro	Asp	Lys	Gln	Met	Ala	Val	Leu	Pro	Arg	Arg	Glu	Arg	Asn	Arg	
1				5					10					15		
Gln	Ala	Ala	Ala	Ala	Asn	Pro	Glu	Asn	Ser	Arg	Gly	Lys	Gly	Arg	Arg	
			20				25					30				
Gly	Gln	Arg	Gly	Lys	Asn	Arg	Gly	Cys	Val	Leu	Thr	Ala	Ile	His	Leu	
		35				40						45				

```

Asn Val Thr Asp Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu Leu Ile
  50                      55                      60
Phe Arg Tyr Cys Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr Tyr Asp
  65                      70                      75                      80
Lys Ile Leu Lys Asn Leu Ser Arg Asn Arg Arg Leu Val Ser Asp Lys
                      85                      90                      95
Val Gly Gln Ala Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp Leu Ser
                      100                      105                      110
Phe Leu Asp Asp Asn Leu Val Tyr His Ile Leu Arg Lys His Ser Ala
                      115                      120                      125
Lys Arg Cys Gly Cys Ile
                      130

```

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 4 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

```

Lys Asn Arg Gly
1

```

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

```

Gly Lys Asn Arg Gly
1                      5

```

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 6 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Arg Gly Lys Asn Arg Gly
1 5

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 7 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Gln Arg Gly Lys Asn Arg Gly
1 5

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Gly Gln Arg Gly Lys Asn Arg Gly
1 5

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 9 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Arg Gly Gln Arg Gly Lys Asn Arg Gly
1 5

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
1 5 10

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
1 5 10

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
1 5 10

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 13 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
1 5 10

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 14 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:
- | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Arg | Gly | Lys | Gly | Arg | Arg | Gly | Gln | Arg | Gly | Lys | Asn | Arg | Gly |
| 1 | | | | 5 | | | | | 10 | | | | |

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:
- | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Ser | Arg | Gly | Lys | Gly | Arg | Arg | Gly | Gln | Arg | Gly | Lys | Asn | Arg | Gly |
| 1 | | | | 5 | | | | | 10 | | | | 15 | |

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 16 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:
- | | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Asn | Ser | Arg | Gly | Lys | Gly | Arg | Arg | Gly | Gln | Arg | Gly | Lys | Asn | Arg | Gly |
| 1 | | | | 5 | | | | | 10 | | | | 15 | | |

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 17 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg
 1 5 10 15
 Gly

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn
 1 5 10 15
 Arg Gly

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys
 1 5 10 15
 Asn Arg Gly

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly
 1 5 10 15
 Lys Asn Arg Gly
 20

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg
 1 5 10 15
 Gly Lys Asn Arg Gly
 20

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln
 1 5 10 15
 Arg Gly Lys Asn Arg Gly
 20

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 23 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly
 1 5 10 15
 Gln Arg Gly Lys Asn Arg Gly
 20

(2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

Gln Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg
 1 5 10 15
 Gly Gln Arg Gly Lys Asn Arg Gly
 20

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 25 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

Arg Gln Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg
 1 5 10 15
 Arg Gly Gln Arg Gly Lys Asn Arg Gly
 20 25

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 26 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Asn Arg Gln Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly
 1 5 10 15

Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
 20 25

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 27 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

Arg Asn Arg Gln Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys
 1 5 10 15

Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
 20 25

(2) INFORMATION FOR SEQ ID NO:27:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 28 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Glu Arg Asn Arg Gln Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly
 1 5 10 15

Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
 20 25

(2) INFORMATION FOR SEQ ID NO:28:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 29 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

Arg Glu Arg Asn Arg Gln Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg
 1 5 10 15

Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
 20 25

(2) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

Arg Arg Glu Arg Asn Arg Gln Ala Ala Ala Asn Pro Glu Asn Ser
 1 5 10 15
 Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
 20 25 30

(2) INFORMATION FOR SEQ ID NO:30:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 31 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Pro Arg Arg Glu Arg Asn Arg Gln Ala Ala Ala Asn Pro Glu Asn
 1 5 10 15
 Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
 20 25 30

(2) INFORMATION FOR SEQ ID NO:31:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

Leu Pro Arg Arg Glu Arg Asn Arg Gln Ala Ala Ala Asn Pro Glu
 1 5 10 15

Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg Gly
 20 25 30

(2) INFORMATION FOR SEQ ID NO:32:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 33 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

Val Leu Pro Arg Arg Glu Arg Asn Arg Gln Ala Ala Ala Ala Asn Pro
 1 5 10 15

Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn Arg
 20 25 30

Gly

(2) INFORMATION FOR SEQ ID NO:33:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 34 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

Ala Val Leu Pro Arg Arg Glu Arg Asn Arg Gln Ala Ala Ala Ala Asn
 1 5 10 15

Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys Asn
 20 25 30

Arg Gly

(2) INFORMATION FOR SEQ ID NO:34:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 35 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

Met Ala Val Leu Pro Arg Arg Glu Arg Asn Arg Gln Ala Ala Ala Ala
 1 5 10 15
 Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly Lys
 20 25 30
 Asn Arg Gly
 35

(2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 36 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

Gln Met Ala Val Leu Pro Arg Arg Glu Arg Asn Arg Gln Ala Ala Ala
 1 5 10 15
 Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly
 20 25 30
 Lys Asn Arg Gly
 35

(2) INFORMATION FOR SEQ ID NO:36:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 37 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

Lys Gln Met Ala Val Leu Pro Arg Arg Glu Arg Asn Arg Gln Ala Ala
 1 5 10 15
 Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg
 20 25 30
 Gly Lys Asn Arg Gly
 35

(2) INFORMATION FOR SEQ ID NO:37:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 38 amino acids

- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

```

Asp Lys Gln Met Ala Val Leu Pro Arg Arg Glu Arg Asn Arg Gln Ala
 1           5           10           15

Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln
          20           25           30

Arg Gly Lys Asn Arg Gly
          35

```

(2) INFORMATION FOR SEQ ID NO:38:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 39 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

```

Pro Asp Lys Gln Met Ala Val Leu Pro Arg Arg Glu Arg Asn Arg Gln
 1           5           10           15

Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly
          20           25           30

Gln Arg Gly Lys Asn Arg Gly
          35

```

(2) INFORMATION FOR SEQ ID NO:39:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 417 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

```

CATATGTCTC CGGATAAACA AATGGCTGTT CTTCCACGTC GTGAACGTAA CCGTCAGGCG 60
GCCGCTGCTA ACCCGGAGAA TTCCCGTGGT AAAGGTCGTC GTGGTCAGCG TGGTAAAAAC 120
CGCGGTTGCG TTCTGACCGC TATCCACCTG AACGTTACCG ACCTGGGTCT CGGTTACGAA 180
ACCAAAGAAG AATTAATCTT CCGTTACTGC TCCGGTTCCT GCGACGCTGC TGAAACCACG 240

```

TACGACAAAA TCCTGAAAAA CCTGTCCCGT AACCGTCGTC TGGTTTCCGA CAAAGTTGGT 300
 CAAGCTTGCT GCGTCCCGAT CGCTTTCGAC GACGACCTGT CCTTCCTGGA CGACAACCTG 360
 GTTTACCACA TCCTGCGTAA AACTCCGCT AAGCGTTGCG GTTGACATCTA AGGATCC 417

(2) INFORMATION FOR SEQ ID NO:40:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 417 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

CATATGAGCC CGGACAAACA GATGGCAGTA CTTCACGTC GTGAACGTAA TCGCCAGGCA 60
 GCAGCTGCAA ACCCGGAAAA CTCCCCTGGT AAAGGTCGCC GTGGCCAGCG CGGCAAAAC 120
 CGTGGTTGTG TTCTGACTGC AATCCACCTG AACGTTACTG ACCTGGGTCT GGGCTACGAA 180
 ACCAAAGAAG AACTGATCTT CCGCTACTGC AGCGGCTCTT GCGACGCAGC TGAAACCACT 240
 TACGACAAAA TCCTGAAAAA CCTGTCCCGT AACCGCCGTC TGGTAAGCGA CAAAGTAGGT 300
 CAGGCATGCT GCGTCCCGAT CGCATTCGAC GATGACCTGA GCTTCCTGGA TGACAACCTG 360
 GTTTACCACA TCCTGCGTAA AACTCCGCT AACCGCTGCG GTTGACATCTA AGGATCC 417

(2) INFORMATION FOR SEQ ID NO:41:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 345 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 1..342

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

ATG TCC CCA GAA AAT TCT CGT GGT AAA GGT CGT CGT GGT CAG CGT GGT 48
 Met Ser Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly
 135 140 145 150
 AAT AAC CGC GGT TGC GTT CTG ACC GCT ATC CAC CTG AAC GTT ACC GAC 96
 Asn Asn Arg Gly Cys Val Leu Thr Ala Ile His Leu Asn Val Thr Asp
 155 160 165

CTG GGT CTC GGT TAC GAA ACC AAA GAA GAA TTA ATC TTC CGT TAC TGC	144
Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu Leu Ile Phe Arg Tyr Cys	
170 175 180	
TCC GGT TCC TGC GAC GCT GCT GAA ACC ACG TAC GAC AAA ATC CTG AAA	192
Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr Tyr Asp Lys Ile Leu Lys	
185 190 195	
AAC CTG TCC CGT AAC CGT CGT CTG GTT TCC GAC AAA GTT GGT CAA GCT	240
Asn Leu Ser Arg Asn Arg Arg Leu Val Ser Asp Lys Val Gly Gln Ala	
200 205 210	
TGC TGC CGT CCG ATC GCT TTC GAC GAC GAC CTG TCC TTC CTG GAC GAC	288
Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp Leu Ser Phe Leu Asp Asp	
215 220 225 230	
AAC CTG GTT TAC CAC ATC CTG CGT AAA CAC TCC GCT AAG CGT TGC GGT	336
Asn Leu Val Tyr His Ile Leu Arg Lys His Ser Ala Lys Arg Cys Gly	
235 240 245	
TGC ATC TAA	345
Cys Ile	

(2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 114 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

Met	Ser	Pro	Glu	Asn	Ser	Arg	Gly	Lys	Gly	Arg	Arg	Gly	Gln	Arg	Gly	1	5	10	15
Asn	Asn	Arg	Gly	Cys	Val	Leu	Thr	Ala	Ile	His	Leu	Asn	Val	Thr	Asp	20	25	30	
Leu	Gly	Leu	Gly	Tyr	Glu	Thr	Lys	Glu	Glu	Leu	Ile	Phe	Arg	Tyr	Cys	35	40	45	
Ser	Gly	Ser	Cys	Asp	Ala	Ala	Glu	Thr	Thr	Tyr	Asp	Lys	Ile	Leu	Lys	50	55	60	
Asn	Leu	Ser	Arg	Asn	Arg	Arg	Leu	Val	Ser	Asp	Lys	Val	Gly	Gln	Ala	65	70	75	80
Cys	Cys	Arg	Pro	Ile	Ala	Phe	Asp	Asp	Asp	Leu	Ser	Phe	Leu	Asp	Asp	85	90	95	
Asn	Leu	Val	Tyr	His	Ile	Leu	Arg	Lys	His	Ser	Ala	Lys	Arg	Cys	Gly	100	105	110	
Cys	Ile																		

(2) INFORMATION FOR SEQ ID NO:43:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 315 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..312

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

ATG	CGT	GGT	CAA	CGT	GGT	AAA	AAC	CGC	GGT	TGC	GTT	CTG	ACT	GCA	ATC	48
Met	Arg	Gly	Gln	Arg	Gly	Lys	Asn	Arg	Gly	Cys	Val	Leu	Thr	Ala	Ile	
115					120					125					130	
CAC	CTG	AAC	GTT	ACT	GAC	CTG	GGT	CTG	GGC	TAC	GAA	ACC	AAA	GAA	GAA	96
His	Leu	Asn	Val	Thr	Asp	Leu	Gly	Leu	Gly	Tyr	Glu	Thr	Lys	Glu	Glu	
				135					140						145	
CTG	ATC	TTC	CGC	TAC	TGC	AGC	GGC	TCT	TGC	GAC	GCA	GCT	GAA	ACC	ACT	144
Leu	Ile	Phe	Arg	Tyr	Cys	Ser	Gly	Ser	Cys	Asp	Ala	Ala	Glu	Thr	Thr	
				150					155					160		
TAC	GAC	AAA	ATC	CTG	AAA	AAC	CTG	TCC	CGT	AAC	CGC	CGT	CTG	GTA	AGC	192
Tyr	Asp	Lys	Ile	Leu	Lys	Asn	Leu	Ser	Arg	Asn	Arg	Arg	Leu	Val	Ser	
				165					170					175		
GAC	AAA	GTA	GGT	CAG	GCA	TGC	TGC	CGT	CCG	ATC	GCA	TTC	GAC	GAT	GAC	240
Asp	Lys	Val	Gly	Gln	Ala	Cys	Cys	Arg	Pro	Ile	Ala	Phe	Asp	Asp	Asp	
				180					185					190		
CTG	AGC	TTC	CTG	GAT	GAC	AAC	CTG	GTT	TAC	CAC	ATC	CTG	CGT	AAA	CAC	288
Leu	Ser	Phe	Leu	Asp	Asp	Asn	Leu	Val	Tyr	His	Ile	Leu	Arg	Lys	His	
				195					200					210		
TCC	GCT	AAA	CGC	TGC	GGT	TGC	ATC	TAA								315
Ser	Ala	Lys	Arg	Cys	Gly	Cys	Ile									
								215								

(2) INFORMATION FOR SEQ ID NO:44:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 104 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

Met	Arg	Gly	Gln	Arg	Gly	Lys	Asn	Arg	Gly	Cys	Val	Leu	Thr	Ala	Ile
1					5				10					15	

His Leu Asn Val Thr Asp Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu
 20 25 30
 Leu Ile Phe Arg Tyr Cys Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr
 35 40 45
 Asp Lys Val Gly Gln Ala Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp
 50 55 60 65 70 75 80
 Leu Ser Phe Leu Asp Asp Asn Leu Val Tyr His Ile Leu Arg Lys His
 85 90 95
 Ser Ala Lys Arg Cys Gly Cys Ile
 100

(2) INFORMATION FOR SEQ ID NO:45:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 312 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 1..309

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

ATG GGT CAA CGT GGT AAA AAC CGT GGT TGT GTT CTG ACT GCA ATC CAC 48
 Met Gly Gln Arg Gly Lys Asn Arg Gly Cys Val Leu Thr Ala Ile His
 105 110 115 120
 CTG AAC GTT ACT GAC CTG GGT CTG GGC TAC GAA ACC AAA GAA GAA CTG 96
 Leu Asn Val Thr Asp Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu Leu
 125 130 135
 ATC TTC CGC TAC TGC AGC GGC TCT TGC GAC GCA GCT GAA ACC ACT TAC 144
 Ile Phe Arg Tyr Cys Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr Tyr
 140 145 150
 GAC AAA ATC CTG AAA AAC CTG TCC CGT AAC CGC CGT CTG GTA AGC GAC 192
 Asp Lys Ile Leu Lys Asn Leu Ser Arg Asn Arg Arg Leu Val Ser Asp
 155 160 165
 AAA GTA GGT CAG GCA TGC TGC CGT CCG ATC GCA TTC GAC GAT GAC CTG 240
 Lys Val Gly Gln Ala Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp Leu
 170 175 180
 AGC TTC CTG GAT GAC AAC CTG GTT TAC CAC ATC CTG CGT AAA CAC TCC 288
 Ser Phe Leu Asp Asp Asn Leu Val Tyr His Ile Leu Arg Lys His Ser
 185 190 195 200

GCT AAA CGC TGC GGT TGC ATC TAA
 Ala Lys Arg Cys Gly Cys Ile
 205

312

(2) INFORMATION FOR SEQ ID NO:46:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 103 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

Met Gly Gln Arg Gly Lys Asn Arg Gly Cys Val Leu Thr Ala Ile His
 1 5 10 15
 Leu Asn Val Thr Asp Leu Gly Leu Gly Tyr Glu Thr Lys Glu Leu
 20 25 30
 Ile Phe Arg Tyr Cys Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr Tyr
 35 40 45
 Asp Lys Ile Leu Lys Asn Leu Ser Arg Asn Arg Arg Leu Val Ser Asp
 50 55 60
 Lys Val Gly Gln Ala Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp Leu
 65 70 75 80
 Ser Phe Leu Asp Asp Asn Leu Val Tyr His Ile Leu Arg Lys His Ser
 85 90 95
 Ala Lys Arg Cys Gly Cys Ile
 100

(2) INFORMATION FOR SEQ ID NO:47:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 135 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

Met Ser Pro Asp Lys Gln Met Ala Val Leu Pro Arg Arg Glu Arg Asn
 1 5 10 15
 Arg Gln Ala Ala Ala Ala Asn Pro Glu Asn Ser Arg Gly Lys Gly Arg
 20 25 30
 Arg Gly Gln Arg Gly Lys Asn Arg Gly Cys Val Leu Thr Ala Ile His
 35 40 45

```

Leu Asn Val Thr Asp Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu Leu
 50                      55                      60

Ile Phe Arg Tyr Cys Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr Tyr
 65                      70                      75                      80

Asp Lys Ile Leu Lys Asn Leu Ser Arg Asn Arg Arg Leu Val Ser Asp
                      85                      90                      95

Lys Val Gly Gln Ala Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp Leu
                      100                     105                     110

Ser Phe Leu Asp Asp Asn Leu Val Tyr His Ile Leu Arg Lys His Ser
                      115                     120                     125

Ala Lys Arg Cys Gly Cys Ile
                      130                     135

```

(2) INFORMATION FOR SEQ ID NO:48:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 104 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

```

Met Arg Gly Gln Arg Gly Lys Asn Arg Gly Cys Val Leu Thr Ala Ile
 1                      5                      10                      15

His Leu Asn Val Thr Asp Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu
                      20                      25                      30

Leu Ile Phe Arg Tyr Cys Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr
                      35                      40                      45

Tyr Asp Lys Ile Leu Lys Asn Leu Ser Arg Asn Arg Arg Leu Val Ser
 50                      55                      60

Asp Lys Val Gly Gln Ala Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp
 65                      70                      75                      80

Leu Ser Phe Leu Asp Asp Asn Leu Val Tyr His Ile Leu Arg Lys His
                      85                      90                      95

Ser Ala Lys Arg Cys Gly Cys Ile
                      100

```

(2) INFORMATION FOR SEQ ID NO:49:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 103 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

```

Met Gly Gln Arg Gly Lys Asn Arg Gly Cys Val Leu Thr Ala Ile His
 1           5           10           15
Leu Asn Val Thr Asp Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu Leu
          20           25           30
Ile Phe Arg Tyr Cys Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr Tyr
          35           40           45
Asp Lys Ile Leu Lys Asn Leu Ser Arg Asn Arg Arg Leu Val Ser Asp
          50           55           60
Lys Val Gly Gln Ala Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp Leu
          65           70           75           80
Ser Phe Leu Asp Asp Asn Leu Val Tyr His Ile Leu Arg Lys His Ser
          85           90           95
Ala Lys Arg Cys Gly Cys Ile
          100

```

(2) INFORMATION FOR SEQ ID NO:50:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 114 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

```

Met Ser Pro Glu Asn Ser Arg Gly Lys Gly Arg Arg Gly Gln Arg Gly
 1           5           10           15
Asn Asn Arg Gly Cys Val Leu Thr Ala Ile His Leu Asn Val Thr Asp
          20           25           30
Leu Gly Leu Gly Tyr Glu Thr Lys Glu Glu Leu Ile Phe Arg Tyr Cys
          35           40           45
Ser Gly Ser Cys Asp Ala Ala Glu Thr Thr Tyr Asp Lys Ile Leu Lys
          50           55           60
Asn Leu Ser Arg Asn Arg Arg Leu Val Ser Asp Lys Val Gly Gln Ala
          65           70           75           80
Cys Cys Arg Pro Ile Ala Phe Asp Asp Asp Leu Ser Phe Leu Asp Asp
          85           90           95

```


A-357

- 82 -

Asn	Leu	Val	Tyr	His	Ile	Leu	Arg	Lys	His	Ser	Ala	Lys	Arg	Cys	Gly
			100					105						110	


Cys Ile

$\frac{2.11}{5.17}$
 $\frac{1.1}{5.07}$
 $\frac{2.09}{6.17}$
 $\frac{2.12}{5.18}$
 $\frac{1.1}{5.18}$
 $\frac{2.11}{5.17}$
 $\frac{2.09}{5.17}$
 $\frac{1.1}{5.20}$
 $\frac{1}{5}$
 $\frac{1}{5.19}$
 $\frac{2.09}{5.17}$
 $\frac{1}{5.21}$
 $\frac{1}{5.12}$
 $\frac{2.11}{5.17}$
 $\frac{2.11}{5.17}$

CLAIMS

What is claimed is:

1. A truncated glial cell line-derived neurotrophic factor (GDNF) protein product having an amino acid sequence



wherein

- [Cys⁴¹-Cys¹³³] represents the amino acid sequence of Cys⁴¹ through Cys¹³³ as depicted in Figure 1 (SEQ ID NO 2);

Y represents the carboxy terminal group of Cys¹³³ or a carboxy-terminus amino acid residue of Ile¹³⁴; and

X represents a methionylated or nonmethionylated amine group of Cys⁴¹ or amino-terminus amino acid residue(s) selected from the group:

15

G
RG
NRG
KNRG (SEQ ID NO:3)
GKNRG (SEQ ID NO:4)
RGKNRG (SEQ ID NO:5)
QRGKNRG (SEQ ID NO:6)
GQRGKNRG (SEQ ID NO:7)
RGQRGKNRG (SEQ ID NO:8)
RRGQRGKNRG (SEQ ID NO:9)
G RRGQRGKNRG (SEQ ID NO:10)
KG RRGQRGKNRG (SEQ ID NO:11)
GKG RRGQRGKNRG (SEQ ID NO:12)
RGKG RRGQRGKNRG (SEQ ID NO:13)
SRGKG RRGQRGKNRG (SEQ ID NO:14)
NSRGKG RRGQRGKNRG (SEQ ID NO:15)
ENSRGKG RRGQRGKNRG (SEQ ID NO:16)
PENSRGKG RRGQRGKNRG (SEQ ID NO:17)
NPENSRGKG RRGQRGKNRG (SEQ ID NO:18)
ANPENSRGKG RRGQRGKNRG (SEQ ID NO:19)
A ANPENSRGKG RRGQRGKNRG (SEQ ID NO:20)

AA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:21)
 AAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:22)
 QAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:23)
 RQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:24)
 NRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:25)
 RNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:26)
 ERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:27)
 RERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:28)
 RRERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:29)
 P RRERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:30)
 LP RRERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:31)
 VLP RRERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:32)
 AVLP RRERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:33)
 MAVLP RRERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:34)
 QMAVLP RRERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:35)
 KQMAVLP RRERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:36)
 DKQMAVLP RRERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:37) and
 PDKQMAVLP RRERNRQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:38)

and addition, substitution and internal deletion variants and derivatives thereof.

2. A truncated GDNF protein product according to Claim 1, wherein X =

5 RQAAA ANPENSRGKG RRGQRGKNRG (SEQ ID NO:24) or a variant thereof.

3. A truncated GDNF protein product according to Claim 1, wherein X =

NPENSRGKG RRGQRGKNRG (SEQ ID NO:18) or a variant thereof.

10 4. A truncated GDNF protein product according to Claim 1, wherein X =

PENSRGKG RRGQRGKNRG (SEQ ID NO:17) or a variant thereof.

5. A truncated GDNF protein product according to Claim 1, wherein X =

SRGKG RRGQRGKNRG (SEQ ID NO:14) or a variant thereof.

15

6. A truncated GDNF protein product according to Claim 1, wherein X =

RGQRGKNRG (SEQ ID NO:8) or a variant thereof.

7. A truncated GDNF protein product according to Claim 1, wherein X = QQRGKNRG (SEQ ID NO:7) or a variant thereof.

8. A truncated GDNF protein product according to Claim 1, wherein X =
5 KNRG (SEQ ID NO:3) or a variant thereof.

9. A truncated GDNF protein product according to Claim 1, wherein X = NRG
or a variant thereof.

10 10. A truncated GDNF protein product according to Claims 1 through 9,
wherein said amino acid sequence is glycosylated.

11. A truncated GDNF protein product according to Claims 1 through 9,
wherein said amino acid sequence is nonglycosylated.

15 12. A truncated GDNF protein product according to Claim 1, wherein said
derivative is an X-[Cys⁴¹-Cys¹³³]-Y amino acid sequence conjugated to a water
soluble polymer.

20 13. A polynucleotide encoding a truncated GDNF protein according to Claim 1.

14. A polynucleotide according to Claim 13, comprising a portion of the
sequence as set forth in Figure 1.

25 15. A polynucleotide according to Claim 13, comprising a portion of the
sequence as set forth in Figure 3.

16. A polynucleotide according to Claim 13, comprising a portion of the
sequence as set forth in Figure 4.

30 17. A polynucleotide according to Claim 13, comprising the sequence as set
forth in Figure 5.

18. A polynucleotide according to Claim 13, comprising the sequence as set
35 forth in Figure 6.

19. A polynucleotide according to Claim 13, comprising the sequence as set forth in Figure 7.

20. A vector, comprising a polynucleotide of Claim 13 operatively linked to an expression control sequence.

21. A prokaryotic or eukaryotic host cell transformed or transfected with a polynucleotide of Claim 13.

22. A method for producing a truncated GDNF protein, comprising growing host cells of Claim 21 in a suitable nutrient medium and, optionally, isolating said truncated GDNF from said cells or said nutrient medium.

23. A method for producing a truncated GDNF protein according to Claim 22, wherein said host cells are *E. coli*.

24. A method for producing a truncated GDNF protein according to Claim 22, wherein said host cells are Chinese hamster ovary cells.

25. A method for the production of a truncated glial cell line-derived neurotrophic factor (GDNF) protein, comprising the steps of:

- (a) culturing a prokaryotic or eukaryotic host cell transformed or transfected with a vector of Claim 20;
- (b) maintaining said host cell under conditions allowing the expression of truncated GDNF protein by said host cell; and
- (c) optionally isolating the truncated GDNF protein expressed by said host cell.

26. A truncated GDNF protein which is the recombinant expression product of a prokaryotic or eukaryotic host cell containing an exogenous polynucleotide of Claim 13.

27. A pharmaceutical composition comprising a truncated GDNF protein product according to Claim 1 in association with a pharmaceutically acceptable vehicle.

28. A pharmaceutical composition comprising a truncated GDNF protein, produced in accordance with the method of Claim 22 in association with a pharmaceutically acceptable vehicle.

5 29. A pharmaceutical composition comprising a truncated GDNF protein, produced in accordance with the method of Claim 25 in association with a pharmaceutically acceptable vehicle.

10 30. A method of treating Parkinson's Disease comprising administering to a patient the pharmaceutical composition of Claim 27.

15 31. A method of treating Parkinson's Disease comprising administering to a patient a polynucleotide sequence of Claim 13 to provide *in vivo* production of said truncated GDNF protein.

32. A method of treating Parkinson's Disease comprising implanting in a patient a cell transformed with a polynucleotide sequence of Claim 13 to provide *in vivo* production of said truncated GDNF protein.

20 33. A glial cell line-derived neurotrophic factor (GDNF) composition, comprising a mature GDNF protein and one or more truncated GDNF proteins, wherein said mature GDNF protein has a molecular weight of approximately 44 kDa, and wherein said truncated GDNF protein(s) has a molecular weight of approximately 29 to 40 kDa.

25 34. A GDNF composition according to Claim 33, comprising at least two species of truncated GDNF protein wherein a first species has a molecular weight of approximately 36 kDa and a second species has a molecular weight of approximately 40 kDa.

30 35. A GDNF composition according to Claim 34, wherein said second truncated GDNF species having a molecular weight of approximately 40 kDa is a heterodimer of a mature GDNF monomer having a molecular weight of approximately 22 kDa and a truncated GDNF monomer having a molecular weight of approximately 18 kDa.

35

36. A truncated GDNF protein isolated from the GDNF composition of Claim 33 and having a molecular weight of approximately 29 to 40 kDa.

37. A truncated GDNF protein isolated from the GDNF composition of Claim 33 and having a molecular weight of approximately 29 to 36 kDa.

38. A truncated GDNF protein derived from a mature GDNF protein expressed by a recombinantly modified bacterial or mammalian cell, said truncated GDNF protein having an amino acid sequence

10 X-[Cys⁴¹-Cys¹³³]-Y

wherein

[Cys⁴¹-Cys¹³³] represents the amino acid sequence of Cys⁴¹ through Cys¹³³ as depicted in Figure 1 (SEQ ID NO 2);

15 Y represents the carboxy terminal group of Cys¹³³ or a carboxy-terminus amino acid residue of Ile¹³⁴; and

X represents an amine group of Cys⁴¹ or amino-terminus amino acid residue(s) selected from the group:

G
RG
NRG
KNRG (SEQ ID NO:3)
GKNRG (SEQ ID NO:4)
RGKNRG (SEQ ID NO:5)
QRGKNRG (SEQ ID NO:6)
GQRGKNRG (SEQ ID NO:7)
RQQRGKNRG (SEQ ID NO:8)
RRQQRGKNRG (SEQ ID NO:9)
G RRGQRGKNRG (SEQ ID NO:10)
KG RRGQRGKNRG (SEQ ID NO:11)
GKG RRGQRGKNRG (SEQ ID NO:12)
RGKG RRGQRGKNRG (SEQ ID NO:13)
SRGKG RRGQRGKNRG (SEQ ID NO:14)
NSRGKG RRGQRGKNRG (SEQ ID NO:15)
ENSRGKG RRGQRGKNRG (SEQ ID NO:16)
PENSRGKG RRGQRGKNRG (SEQ ID NO:17)

		NPENSRGKG	RRGQRGKNRG	(SEQ ID NO:18)
		ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:19)
	A	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:20)
	AA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:21)
	AAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:22)
	QAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:23)
	RQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:24)
	NRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:25)
	RNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:26)
	ERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:27)
	RERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:28)
	RRERNRQAAA	ANPENSRGKG	RRGQRGKNRG	(SEQ ID NO:29)
	P	RRERNRQAAA	ANPENSRGKG	(SEQ ID NO:30)
	LP	RRERNRQAAA	ANPENSRGKG	(SEQ ID NO:31)
	VLP	RRERNRQAAA	ANPENSRGKG	(SEQ ID NO:32)
	AVLP	RRERNRQAAA	ANPENSRGKG	(SEQ ID NO:33)
	MAVLP	RRERNRQAAA	ANPENSRGKG	(SEQ ID NO:34)
	QMAVLP	RRERNRQAAA	ANPENSRGKG	(SEQ ID NO:35)
	KQMAVLP	RRERNRQAAA	ANPENSRGKG	(SEQ ID NO:36)
	DKQMAVLP	RRERNRQAAA	ANPENSRGKG	(SEQ ID NO:37) and
	PDKQMAVLP	RRERNRQAAA	ANPENSRGKG	(SEQ ID NO:38)

and addition, substitution and internal deletion variants thereof.

39. A truncated GDNF protein according to Claim 38, wherein X is selected
 5 from the group consisting of:

G
 RG
 NRG
 KNRG (SEQ ID NO:3)
 GKNRG (SEQ ID NO:4)
 RGKNRG (SEQ ID NO:5)
 QRGKNRG (SEQ ID NO:6)
 GQRGKNRG (SEQ ID NO:7)
 RGQRGKNRG (SEQ ID NO:8) and
 RRGQRGKNRG (SEQ ID NO:9)

and variants thereof.

40. A truncated GDNF protein according to Claim 38, wherein mature GDNF protein is expressed by a recombinantly modified bacterial cell and said
5 truncated GDNF protein is produced *in vitro* or *in vivo*.

41. A method of preparing a pharmaceutical composition wherein a therapeutically effective amount of a truncated GDNF protein product according to Claim 1 is mixed with one or more pharmaceutically acceptable vehicles.
10

42. The use of a truncated GDNF protein product according to Claim 1 for treating damage to the nervous system caused by disease or injury.

43. The use of a truncated GDNF protein product according to Claim 42 for
15 treating Parkinson's disease.

44. The use of a truncated GDNF protein product according to Claim 1 for the preparation of a pharmaceutical composition for treating damage to the nervous system caused by disease or injury.
20

TRUNCATED GLIAL CELL LINE-DERIVED NEUROTROPHIC FACTOR

ABSTRACT

- 5 Disclosed are novel proteins, referred to as truncated glial cell line-derived neurotrophic factor (truncated GDNF) proteins, that promote dopamine uptake by dopaminergic cells and promote the survival of nerve cells. Also disclosed are processes for obtaining the truncated GDNF proteins by recombinant genetic engineering techniques.

10

11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 100
 101
 102
 103
 104
 105
 106
 107
 108
 109
 110
 111
 112
 113
 114
 115
 116
 117
 118
 119
 120
 121
 122
 123
 124
 125
 126
 127
 128
 129
 130
 131
 132
 133
 134
 135
 136
 137
 138
 139
 140
 141
 142
 143
 144
 145
 146
 147
 148
 149
 150
 151
 152
 153
 154
 155
 156
 157
 158
 159
 160
 161
 162
 163
 164
 165
 166
 167
 168
 169
 170
 171
 172
 173
 174
 175
 176
 177
 178
 179
 180
 181
 182
 183
 184
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201
 202
 203
 204
 205
 206
 207
 208
 209
 210
 211
 212
 213
 214
 215
 216
 217
 218
 219
 220
 221
 222
 223
 224
 225
 226
 227
 228
 229
 230
 231
 232
 233
 234
 235
 236
 237
 238
 239
 240
 241
 242
 243
 244
 245
 246
 247
 248
 249
 250
 251
 252
 253
 254
 255
 256
 257
 258
 259
 260
 261
 262
 263
 264
 265
 266
 267
 268
 269
 270
 271
 272
 273
 274
 275
 276
 277
 278
 279
 280
 281
 282
 283
 284
 285
 286
 287
 288
 289
 290
 291
 292
 293
 294
 295
 296
 297
 298
 299
 300
 301
 302
 303
 304
 305
 306
 307
 308
 309
 310
 311
 312
 313
 314
 315
 316
 317
 318
 319
 320
 321
 322
 323
 324
 325
 326
 327
 328
 329
 330
 331
 332
 333
 334
 335
 336
 337
 338
 339
 340
 341
 342
 343
 344
 345
 346
 347
 348
 349
 350
 351
 352
 353
 354
 355
 356
 357
 358
 359
 360
 361
 362
 363
 364
 365
 366
 367
 368
 369
 370
 371
 372
 373
 374
 375
 376
 377
 378
 379
 380
 381
 382
 383
 384
 385
 386
 387
 388
 389
 390
 391
 392
 393
 394
 395
 396
 397
 398
 399
 400
 401
 402
 403
 404
 405
 406
 407
 408
 409
 410
 411
 412
 413
 414
 415
 416
 417
 418
 419
 420
 421
 422
 423
 424
 425
 426
 427
 428
 429
 430
 431
 432
 433
 434
 435
 436
 437
 438
 439
 440
 441
 442
 443
 444
 445
 446
 447
 448
 449
 450
 451
 452
 453
 454
 455
 456
 457
 458
 459
 460
 461
 462
 463
 464
 465
 466
 467
 468
 469
 470
 471
 472
 473
 474
 475
 476
 477
 478
 479
 480
 481
 482
 483
 484
 485
 486
 487
 488
 489
 490
 491
 492
 493
 494
 495
 496
 497
 498
 499
 500
 501
 502
 503
 504
 505
 506
 507
 508
 509
 510
 511
 512
 513
 514
 515
 516
 517
 518
 519
 520
 521
 522
 523
 524
 525
 526
 527
 528
 529
 530
 531
 532
 533
 534
 535
 536
 537
 538
 539
 540
 541
 542
 543
 544
 545
 546
 547
 548
 549
 550
 551
 552
 553
 554
 555
 556
 557
 558
 559
 560
 561
 562
 563
 564
 565
 566
 567
 568
 569
 570
 571
 572
 573
 574
 575
 576
 577
 578
 579
 580
 581
 582
 583
 584
 585
 586
 587
 588
 589
 590
 591
 592
 593
 594
 595
 596
 597
 598
 599
 600
 601
 602
 603
 604
 605
 606
 607
 608
 609
 610
 611
 612
 613
 614
 615
 616
 617
 618
 619
 620
 621
 622
 623
 624
 625
 626
 627
 628
 629
 630
 631
 632
 633
 634
 635
 636
 637
 638
 639
 640
 641
 642
 643
 644
 645
 646
 647
 648
 649
 650
 651
 652
 653
 654
 655
 656
 657
 658
 659
 660
 661
 662
 663
 664
 665
 666
 667
 668
 669
 670
 671
 672
 673
 674
 675
 676
 677
 678
 679
 680
 681
 682
 683
 684
 685
 686
 687
 688
 689
 690
 691
 692
 693
 694
 695
 696
 697
 698
 699
 700
 701
 702
 703
 704
 705
 706
 707
 708
 709
 710
 711
 712
 713
 714
 715
 716
 717
 718
 719
 720
 721
 722
 723
 724
 725
 726
 727
 728
 729
 730
 731
 732
 733
 734
 735
 736
 737
 738
 739
 740
 741
 742
 743
 744
 745
 746
 747
 748
 749
 750
 751
 752
 753
 754
 755
 756
 757
 758
 759
 760
 761
 762
 763
 764
 765
 766
 767
 768
 769
 770
 771
 772
 773
 774
 775
 776
 777
 778
 779
 780
 781
 782
 783
 784
 785
 786
 787
 788
 789
 790
 791
 792
 793
 794
 795
 796
 797
 798
 799
 800
 801
 802
 803
 804
 805
 806
 807
 808
 809
 810
 811
 812
 813
 814
 815
 816
 817
 818
 819
 820
 821
 822
 823
 824
 825
 826
 827
 828
 829
 830
 831
 832
 833
 834
 835
 836
 837
 838
 839
 840
 841
 842
 843
 844
 845
 846
 847
 848
 849
 850
 851
 852
 853
 854
 855
 856
 857
 858
 859
 860
 861
 862
 863
 864
 865
 866
 867
 868
 869
 870
 871
 872
 873
 874
 875
 876
 877
 878
 879
 880
 881
 882
 883
 884
 885
 886
 887
 888
 889
 890
 891
 892
 893
 894
 895
 896
 897
 898
 899
 900
 901
 902
 903
 904
 905
 906
 907
 908
 909
 910
 911
 912
 913
 914
 915
 916
 917
 918
 919
 920
 921
 922
 923
 924
 925
 926
 927
 928
 929
 930
 931
 932
 933
 934
 935
 936
 937
 938
 939
 940
 941
 942
 943
 944
 945
 946
 947
 948
 949
 950
 951
 952
 953
 954
 955
 956
 957
 958
 959
 960
 961
 962
 963
 964
 965
 966
 967
 968
 969
 970
 971
 972
 973
 974
 975
 976
 977
 978
 979
 980
 981
 982
 983
 984
 985
 986
 987
 988
 989
 990
 991
 992
 993
 994
 995
 996
 997
 998
 999
 1000
 1001
 1002
 1003
 1004
 1005
 1006
 1007
 1008
 1009
 1010
 1011
 1012
 1013
 1014
 1015
 1016
 1017
 1018
 1019
 1020
 1021
 1022
 1023
 1024
 1025
 1026
 1027
 1028
 1029
 1030
 1031
 1032
 1033
 1034
 1035
 1036
 1037
 1038
 1039
 1040
 1041
 1042
 1043
 1044
 1045
 1046
 1047
 1048
 1049
 1050
 1051
 1052
 1053
 1054
 1055
 1056
 1057
 1058
 1059
 1060
 1061
 1062
 1063
 1064
 1065
 1066
 1067
 1068
 1069
 1070
 1071
 1072
 1073
 1074
 1075
 1076
 1077
 1078
 1079
 1080
 1081
 1082
 1083
 1084
 1085
 1086
 1087
 1088
 1089
 1090
 1091
 1092
 1093
 1094
 1095
 1096
 1097
 1098
 1099
 1100
 1101
 1102
 1103
 1104
 1105
 1106
 1107
 1108
 1109
 1110
 1111
 1112
 1113
 1114
 1115
 1116
 1117
 1118
 1119
 1120
 1121
 1122
 1123
 1124
 1125
 1126
 1127
 1128
 1129
 1130
 1131
 1132
 1133
 1134
 1135
 1136
 1137
 1138
 1139
 1140
 1141
 1142
 1143
 1144
 1145
 1146
 1147
 1148
 1149
 1150
 1151
 1152
 1153
 1154
 1155
 1156
 1157
 1158
 1159
 1160
 1161
 1162
 1163
 1164
 1165
 1166
 1167
 1168
 1169
 1170
 1171
 1172
 1173
 1174
 1175
 1176
 1177
 1178
 1179
 1180
 1181
 1182
 1183
 1184
 1185
 1186
 1187
 1188
 1189
 1190
 1191
 1192
 1193
 1194
 1195
 1196
 1197
 1198
 1199
 1200
 1201
 1202
 1203
 1204
 1205
 1206
 1207
 1208
 1209
 1210
 1211
 1212
 1213
 1214
 1215
 1216
 1217
 1218
 1219
 1220
 1221
 1222
 1223
 1224
 1225
 1226
 1227
 1228
 1229
 1230
 1231
 1232
 1233
 1234
 1235
 1236
 1237
 1238
 1239
 1240
 1241
 1242
 1243
 1244
 1245
 1246
 1247
 1248
 1249
 1250
 1251
 1252
 1253
 1254
 1255
 1256
 1257
 1258
 1259
 1260
 1261
 1262
 1263
 1264
 1265
 1266
 1267
 1268
 1269
 1270
 1271
 1272
 1273
 1274
 1275
 1276
 1277
 1278
 1279
 1280
 1281
 1282
 1283
 1284
 1285
 1286
 1287
 1288
 1289
 1290
 1291
 1292
 1293
 1294
 1295
 1296
 1297
 1298
 1299
 1300
 1301
 1302
 1303
 1304
 1305
 1306
 1307
 1308
 1309
 1310
 1311
 1312
 1313
 1314
 1315
 1316
 1317
 1318
 1319
 1320
 1321
 1322
 1323
 1324
 1325
 1326
 1327
 1328
 1329
 1330
 1331
 1332
 1333
 1334
 1335
 1336
 1337
 1338
 1339
 1340
 1341
 1342
 1343
 1344
 1345
 1346
 1347
 1348
 1349
 1350
 1351
 1352
 1353
 1354
 1355
 1356
 1357
 1358
 1359
 1360
 1361
 1362
 1363
 1364
 1365
 1366
 1367
 1368
 1369
 1370
 1371
 1372
 1373
 1374
 1375
 1376
 1377
 1378
 1379
 1380
 1381
 1382
 1383
 1384
 1385
 1386
 1387
 1388
 1389
 1390
 1391
 1392
 1393
 1394
 1395
 1396
 1397
 1398
 1399
 1400
 1401
 1402
 1403
 1404
 1405
 1406
 1407
 1408
 1409
 1410
 1411
 1412
 1413
 1414
 1415
 1416
 1417
 1418
 1419
 1420
 1421
 1422
 1423
 1424
 1425
 1426
 1427
 1428
 1429
 1430
 1431
 1432
 1433
 1434
 1435
 1436
 1437
 1438
 1439
 1440
 1441
 1442
 1443
 1444
 1445
 1446
 1447
 1448
 1449
 1450
 1451
 1452
 1453
 1454
 1455
 1456
 1457
 1458
 1459
 1460
 1461
 1462
 1463
 1464
 1465
 1466
 1467
 146

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{n} \log \frac{1}{n} \sum_{i=1}^n \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2} \left(\frac{y_i - \mu}{\sigma} \right)^2} &= \lim_{n \rightarrow \infty} \frac{1}{n} \log \frac{1}{n} \sum_{i=1}^n \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2} \left(\frac{y_i - \mu}{\sigma} \right)^2} \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \log \frac{1}{n} \sum_{i=1}^n \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2} \left(\frac{y_i - \mu}{\sigma} \right)^2} \end{aligned}$$
[illegible]

TCA Ser	CCA Pro	GAT Asp	AAA Lys	CAA Gln 5	ATG Met	GCA Ala	GTG Val	CTT Leu	CCT Pro 10	AGA Arg	AGA Arg	GAG Glu	CGG Arg	AAT Asn 15
CGG Arg	CAG Gln	GCT Ala	GCA Ala	GCT Ala 20	GCC Ala	AAC Asn	CCA Pro	GAG Glu	AAT Asn 25	TCC Ser	AGA Arg	GGA Gly	AAA Lys	GGT Gly 30
CGG Arg	AGA Arg	GGC Gly	CAG Gln	AGG Arg 35	GGC Gly	AAA Lys	AAC Asn	CGG Arg	GGT Gly 40	TGT Cys	GTC Val	TTA Leu	ACT Thr	GCA Ala 45
ATA Ile	CAT His	TTA Leu	AAT Asn	GTC Val 50	ACT Thr	GAC Asp	TTG Leu	GGT Gly	CTG Leu 55	GGC Gly	TAT Tyr	GAA Glu	ACC Thr	AAG Lys 60
GAG Glu	GAA Glu	CTG Leu	ATT Ile	TTT Phe 65	AGG Arg	TAC Tyr	TGC Cys	AGC Ser	GGC Gly 70	TCT Ser	TGC Cys	GAT Asp	GCA Ala	GCT Ala 75
GAG Glu	ACA Thr	ACG Thr	TAC Tyr	GAC Asp 80	AAA Lys	ATA Ile	TTG Leu	AAA Lys	AAC Asn 85	TTA Leu	TCC Ser	AGA Arg	AAT Asn	AGA Arg 90
AGG Arg	CTG Leu	GTG Val	AGT Ser	GAC Asp 95	AAA Lys	GTA Val	GGG Gly	CAG Gln	GCA Ala 100	TGT Cys	TGC Cys	AGA Arg	CCC Pro	ATC Ile 105
GCC Ala	TTT Phe	GAT Asp	GAT Asp	GAC Asp 110	CTG Leu	TCG Ser	TTT Phe	TTA Leu	GAT Asp 115	GAT Asp	AAC Asn	CTG Leu	GTT Val	TAC Tyr 120
CAT His	ATT Ile	CTA Leu	AGA Arg	AAG Lys 125	CAT His	TCC Ser	GCT Ala	AAA Lys	AGG Arg 130	TGT Cys	GGA Gly	TGT Cys	ATC Ile	

FIG. 2




FIG.3A

metGDNF Degenerate DNA Sequence




FIG.3B

S
u
n
I

241 CCTGCGACGCTGCTGAAACCACGTACGACAAAATCCTGAAAAACCTGTCCCGTAACCGTC 300

H E
i a
n m
d l
I P
I v
I u
I I

301 GTCTGGTTTCCGACAAAAGTTGGTCAAGCTTGCTGCCGTC CGATCGCTTTCGACGACGACC 360

361 TGTCTTCCTGGACGACAACCTGTTTACCACATCCTGCGTAAACACTCCGCTAAGCGTT 420

B
a
m
H
I

421 GCGGTTGCATCTAAGGATCC 440

metGDNF Degenerate DNA Sequence

$$\frac{\partial^2 \mathcal{L}}{\partial \theta^2} = \frac{\partial^2 \mathcal{L}}{\partial \theta^2} \bigg|_{\theta = \theta^*} + \frac{\partial^2 \mathcal{L}}{\partial \theta^2} \bigg|_{\theta = \theta^*} \frac{\partial \theta^*}{\partial \theta} + \frac{\partial^2 \mathcal{L}}{\partial \theta^2} \bigg|_{\theta = \theta^*} \frac{\partial^2 \theta^*}{\partial \theta^2} + \dots$$

FIG.5

[Pro²³-Lys³⁷ΔAsn³⁷-Ile¹³⁴] Truncated GDNF Protein

```

ATGTCCCAGAAAATTCTCGTGGTAAAGGTCGTCGGTCAGCGTGGTAATAACCGCGGT
21 -----+-----+-----+-----+-----+-----+ 80
  M S P E N S R G K G R R G Q R G N N R G

TGC GTTCTGACCGCTATCCACCTGAACGTTACCGACCTGGGTCTCGGTACGAAACCAA
81 -----+-----+-----+-----+-----+-----+ 140
  C V L T A I H L N V T D L G L G Y E T K

GAAGAATTAATCTTCCGTTACTGCTCCGGTTCCTGCGACGCTGCTGAAACCACGTACGAC
141 -----+-----+-----+-----+-----+-----+ 200
  E E L I F R Y C S G S C D A A E T T Y D

AAATCCTGAAAAACCTGTCCCGTAACCGTCGTCGGTTTCCGACAAAGTTGGTCAAGCT
201 -----+-----+-----+-----+-----+-----+ 260
  K I L K N L S R N R R L V S D K V G Q A

TGCTGCCGTCCGATCGCTTTCGACGACGACCTGTCCTTCCTGGACGACAACCTGGTTTAC
261 -----+-----+-----+-----+-----+-----+ 320
  C C R P I A F D D D L S F L D D N L V Y

CACATCTGCGTAAACACTCCGCTAAGCGTTGCGGTTGCATCTAA
321 -----+-----+-----+-----+-----+-----+
  H I L R K H S A K R C G C I *

```


FIG.6

[Arg³²-Ile¹³⁴] Truncated GDNF Protein

```

ATGCGTGGTCAACGTGGTAAAAACCGCGTTGCGTTCTGACTGCAATCCACCTGAACGTT
41 -----+-----+-----+-----+-----+-----+ 100
M R G Q R G K N R G C V L T A I H L N V

ACTGACCTGGGTCTGGGCTACGAAACCAAAGAAGAACTGATCTTCCGCTACTGCAGCGGC
101 -----+-----+-----+-----+-----+-----+ 160
T D L G L G Y E T K E E L I F R Y C S G

TCTTTCGACGCAGCTGAAACCACTTACGACAAAATCCTGAAAAACCTGTCCCGTAACCGC
161 -----+-----+-----+-----+-----+-----+ 220
S C D A A E T T Y D K I L K N L S R N R

CGTCTGGTAAGCGACAAAGTAGGTCAGGCATGCTGCCGTCCGATCGCATTCGACGATGAC
221 -----+-----+-----+-----+-----+-----+ 280
R L V S D K V G Q A C C R P I A F D D D

CTGAGCTTCCTGGATGACAACCTGGTTTACCACATCCTGCGTAAACACTCCGCTAAACGC
281 -----+-----+-----+-----+-----+-----+ 340
L S F L D D N L V Y H I L R K H S A K R

TGCGGTTGCATCTAA
341 -----+----- 355
C G C I *
```

FIG.7

[Gly³³-Ile¹³⁴] Truncated GDNF Protein

```

ATGGGTCAACGTGGTAAAAACCGTGGTTGTGTTCTGACTGCAATCCACCTGAACGTTACT
41  -----+-----+-----+-----+-----+-----+ 100
   M  G  Q  R  G  K  N  R  G  C  V  L  T  A  I  H  L  N  V  T

GACCTGGGTCTGGGCTACGAAACCAAAGAAGAACTGATCTTCCGCTACTGCAGCGGCTCT
101 -----+-----+-----+-----+-----+-----+ 160
   D  L  G  L  G  Y  E  T  K  E  E  L  I  F  R  Y  C  S  G  S

TGCAGCGAGCTGAAACCACCTTACGACAAAATCCTGAAAAACCTGTCCCGTAACCGCCGT
161 -----+-----+-----+-----+-----+-----+ 220
   C  D  A  A  E  T  T  Y  D  K  I  L  K  N  L  S  R  N  R  R

CTGGTAAGCGACAAAGTAGGTCAGGCATGCTGCCGTCCGATCGCATTCGACGATGACCTG
221 -----+-----+-----+-----+-----+-----+ 280
   L  V  S  D  K  V  G  Q  A  C  C  R  P  I  A  F  D  D  D  L

AGCTTCTCGGATGACAACCTGGTTTACCACATCCTGCGTAAACACTCCGTAAACGCTGC
281 -----+-----+-----+-----+-----+-----+ 340
   S  F  L  D  D  N  L  V  Y  H  I  L  R  K  H  S  A  K  R  C

GGTTGCATCTAA
341 -----+--- 352
   G  C  I  *
  
```

FIG.8

Comparison of Protein Sequences

					50
	GDNF	MSPDKQMAVL	PRRERNQAA	AANPENSRGK	GRRGQRGKNR GCVLTAIHLN
-31	GDNFMRGQRGKNR GCVLTAIHLN
-32	GDNFMRGQRGKNR GCVLTAIHLN
-22	GDNFMSPENSRGK	GRRGQRGNR GCVLTAIHLN
		51			100
	GDNF	VTDLGLGYET	KEELIFRYCS	GSCDAAETTY	DKILKNLSRN RRLVSDKVGQ
-31	GDNF	VTDLGLGYET	KEELIFRYCS	GSCDAAETTY	DKILKNLSRN RRLVSDKVGQ
-32	GDNF	VTDLGLGYET	KEELIFRYCS	GSCDAAETTY	DKILKNLSRN RRLVSDKVGQ
-22	GDNF	VTDLGLGYET	KEELIFRYCS	GSCDAAETTY	DKILKNLSRN RRLVSDKVGQ
		101		135	
	GDNF	ACCRPIAFDD	DLSFLDDNLV	YHILRKHS AK	RCGCI
-31	GDNF	ACCRPIAFDD	DLSFLDDNLV	YHILRKHS AK	RCGCI
-32	GDNF	ACCRPIAFDD	DLSFLDDNLV	YHILRKHS AK	RCGCI
-22	GDNF	ACCRPIAFDD	DLSFLDDNLV	YHILRKHS AK	RCGCI

DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first, and sole inventor (if only one name is listed below) or a joint inventor (if plural names are listed below) of the invention entitled:

TRUNCATED GLIAL CELL LINE-DERIVED NEUROTROPHIC FACTOR

which is described and claimed in the specification which:

☒ is attached hereto.
☐ was filed on _____
 as Application Serial No.: _____
 and was amended on _____ (if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a).

Power of Attorney: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith:

Ron K. Levy, Registration No.: 31,539, Steven M. Odre, Registration No.: 29,094, and Daniel R. Curry, Registration No. 32,727, said attorney(s)/agent(s) to have in addition full power of revocation, including the power to revoke any power herein granted.

Please send all future correspondence to:

U.S. Patent Operations/DRC
 M/S 10-1-B
 AMGEN INC.
 Amgen Center
 1840 Dehavilland Drive
 Thousand Oaks, California 91320-1789

Direct Telephone Calls To:

Daniel R. Curry
 Attorney/Agent for Applicant(s)
 Registration No.: 32,727
 Phone: (805) 447-8102
 Date: September 28, 1995

Express Mail [®] mail labeling number: YB 530269575 US	Date of Deposit: September 28, 1995
I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. 1.10 on the date indicated above and is addressed to Box Patent Application, Assistant Commissioner of Patents, Washington, D. C. 20231 (Senders Signature Below)	
(Typed name) Ronda G. Spahr	<i>Ronda G Spahr</i>

DECLARATION AND POWER OF ATTORNEY (cont'd)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Full Name of Sole
or First Inventor:

Shaw-Fen Sylvia Hu

Inventor's Signature:

Shaw-Fen Sylvia Hu

Date: Sept 28, 1995

Residence:

986 Lynnmore Drive

Post Office Address:

Thousand Oaks, California 91360 U.S.A.

Citizenship:

U.S.A.

Full Name of Second
Joint Inventor, if Any:

Inventor's Signature:

Date: _____

Residence:

Post Office Address:

Citizenship:

Full Name of Third
Joint Inventor, if Any:

Inventor's Signature:

Date: _____

Residence:

Post Office Address:

Citizenship:
